

IL-27 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1707b**Specification**

IL-27 Antibody (Center) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	Q8TAD2
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	21893
Antigen Region	57-87

IL-27 Antibody (Center) - Additional Information**Gene ID** 53342**Other Names**

Interleukin-17D, IL-17D, Interleukin-27, IL-27, IL17D, IL27

Target/Specificity

This IL-27 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 57-87 amino acids from the Central region of human IL-27.

Dilution

WB~~1:1000
IHC-P~~1:10~50
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IL-27 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

IL-27 Antibody (Center) - Protein Information**Name** IL17D**Function** Induces expression of IL6, CXCL8/IL8, and CSF2/GM-CSF from endothelial cells.

Cellular Location
Secreted.


Tissue Location
Expressed preferentially in adipose, skeletal muscle and CNS.

IL-27 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


IL-27 Antibody (Center) - Images



Western blot analysis of IL27 (arrow) using rabbit polyclonal IL27 Antibody (C71) (Cat. #AP1707b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the IL27 gene.



IL-27 Antibody (Center) (Cat. #AP1707b) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of IL-27 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



IL-27 Antibody (Center) (Cat. #AP1707b) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

IL-27 Antibody (Center) - Background

IL-27 is a cytokine that shares the sequence similarity with IL17. The treatment of endothelial cells with this cytokine has been shown to stimulate the production of other cytokines including IL6, IL8 and CSF2/ GM-CSF. The increased expression of IL8 induced by this cytokine was found to be NF-kappa B-dependent.

IL-27 Antibody (Center) - References

Clark, H.F., et al., Genome Res. 13(10):2265-2270 (2003).
Starnes, T., et al., J. Immunol. 169(2):642-646 (2002).