

IL29 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1709B**Specification**

IL29 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q8IU54
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	54-85

IL29 Antibody (Center) - Additional Information**Gene ID** 282618**Other Names**

Interferon lambda-1, IFN-lambda-1, Cytokine Zcyto21, Interleukin-29, IL-29, IFNL1, IL29, ZCYTO21

Target/Specificity

This IL29 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 54-85 amino acids from the Central region of human IL29.

Dilution

WB~~1:1000

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IL29 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

IL29 Antibody (Center) - Protein Information**Name** IFNL1**Synonyms** IL29, ZCYTO21**Function** Cytokine with antiviral, antitumour and immunomodulatory activities. Plays a critical

role in the antiviral host defense, predominantly in the epithelial tissues. Acts as a ligand for the heterodimeric class II cytokine receptor composed of IL10RB and IFNLR1, and receptor engagement leads to the activation of the JAK/STAT signaling pathway resulting in the expression of IFN-stimulated genes (ISG), which mediate the antiviral state. Has a restricted receptor distribution and therefore restricted targets: is primarily active in epithelial cells and this cell type-selective action is because of the epithelial cell-specific expression of its receptor IFNLR1. Exerts an immunomodulatory effect by up-regulating MHC class I antigen expression.

Cellular Location


Secreted.

IL29 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IL29 Antibody (Center) - Images



The anti-IL29 Pab (Cat. #AP1709b) is used in Western blot to detect IL29 in Jurkat cell lysate.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

IL29 Antibody (Center) - Background

IL29 is a cytokine distantly related to type I interferons and the IL-10 family. This gene, interleukin 28A (IL28A), and interleukin 28B (IL28B) are three closely related cytokine genes that form a cytokine gene cluster on a chromosomal region mapped to 19q13. Expression of the cytokines encoded by the three genes can be induced by viral infection. All three cytokines have been shown to interact with a heterodimeric class II cytokine receptor that consists of interleukin 10 receptor, beta (IL10RB) and interleukin 28 receptor, alpha (IL28RA).

IL29 Antibody (Center) - References

Kotenko, S.V., et al., Nat. Immunol. 4(1):69-77 (2003). Sheppard, P., et al., Nat. Immunol. 4(1):63-68 (2003).