

## ASF1A Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP17195B

## Specification

# ASF1A Antibody (C-term) - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Isotype Calculated MW Antigen Region WB,E <u>O9Y294</u> <u>O3C1E9</u>, <u>O9COE6</u>, <u>O2KIG1</u>, <u>NP\_054753.1</u> Human Bovine, Mouse, Chicken Rabbit Polyclonal Rabbit IgG 22969 140-168

## ASF1A Antibody (C-term) - Additional Information

Gene ID 25842

**Other Names** Histone chaperone ASF1A, Anti-silencing function protein 1 homolog A, hAsf1, hAsf1a, CCG1-interacting factor A, CIA, hCIA, ASF1A

## Target/Specificity

This ASF1A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 140-168 amino acids from the C-terminal region of human ASF1A.

Dilution WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

ASF1A Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

# ASF1A Antibody (C-term) - Protein Information

Name ASF1A {ECO:0000303|PubMed:10759893, ECO:0000312|HGNC:HGNC:20995}



**Function** Histone chaperone that facilitates histone deposition and histone exchange and removal during nucleosome assembly and disassembly (PubMed:<u>10759893</u>, PubMed:<u>11897662</u>, PubMed:<u>12842904</u>, PubMed:<u>14718166</u>, PubMed:<u>15664198</u>, PubMed:<u>16151251</u>, PubMed:<u>21454524</u>). Cooperates with chromatin assembly factor 1 (CAF-1) to promote replication-dependent chromatin assembly and with HIRA to promote replication-independent chromatin assembly (PubMed:<u>11897662</u>, PubMed:<u>14718166</u>, PubMed:<u>15664198</u>). Promotes homologous recombination-mediated repair of double-strand breaks (DSBs) at stalled or collapsed replication forks: acts by mediating histone replacement at DSBs, leading to recruitment of the MMS22L-TONSL complex and subsequent loading of RAD51 (PubMed:<u>29478807</u>). Also involved in the nuclear import of t

Cellular Location Nucleus. Chromosome

**Tissue Location** Ubiquitously expressed.

# ASF1A Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

ASF1A Antibody (C-term) - Images

Hela 55 36 -28 17

ASF1A Antibody (C-term) (Cat. #AP17195b) western blot analysis in Hela cell line lysates (35ug/lane). This demonstrates the ASF1A antibody detected the ASF1A protein (arrow).



# ASF1A Antibody (C-term) - Background

This gene encodes a member of the H3/H4 family of histone chaperone proteins and is similar to the anti-silencing function-1 gene in yeast. The protein is a key component of a histone donor complex that functions in nucleosome assembly. It interacts with histones H3 and H4, and functions together with a chromatin assembly factor during DNA replication and repair. [provided by RefSeq].

## ASF1A Antibody (C-term) - References

Jasencakova, Z., et al. Mol. Cell 37(5):736-743(2010) Newton-Cheh, C., et al. Nat. Genet. 41(4):399-406(2009) Banumathy, G., et al. Mol. Cell. Biol. 29(3):758-770(2009) Ambagala, A.P., et al. J. Virol. 83(1):200-209(2009) Pilyugin, M., et al. PLoS ONE 4 (12), E8328 (2009) :