10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

UGT2B10 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17222B

Specification

UGT2B10 Antibody (C-term) - Product Information

Application	**WB,E**
Primary Accession	P36537
Other Accession	O97951, O75310, O02663, P16662, NP_001066.1
Reactivity	**Human**
Predicted	**Monkey**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**60774**
Antigen Region	**431-458**

UGT2B10 Antibody (C-term) - Additional Information

Gene ID 7365

Other Names
UDP-glucuronosyltransferase 2B10, UDPGT 2B10, UGT2B10

Target/Specificity
This UGT2B10 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 431-458 amino acids from the C-terminal region of human UGT2B10.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
UGT2B10 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

UGT2B10 Antibody (C-term) - Protein Information

Name UGT2B10

Function UDPGT is of major importance in the conjugation and subsequent elimination of potentially toxic xenobiotics and endogenous compounds.


Cellular Location
Microsome membrane; Single-pass membrane protein. Endoplasmic reticulum membrane; Single-pass membrane protein

UGT2B10 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

UGT2B10 Antibody (C-term) - Images



UGT2B10 Antibody (C-term) (Cat. #AP17222b) western blot analysis in 293 cell line lysates (35ug/lane).This demonstrates the UGT2B10 antibody detected the UGT2B10 protein (arrow).

UGT2B10 Antibody (C-term) - Background

UDPGT is of major importance in the conjugation and subsequent elimination of potentially toxic xenobiotics and endogenous compounds.

UGT2B10 Antibody (C-term) - References

Chen, G., et al. Cancer Res. 70(19):7543-7552(2010)
Berg, J.Z., et al. Cancer Epidemiol. Biomarkers Prev. 19(6):1423-1431(2010)
Chen, G., et al. Pharmacogenet. Genomics 18(3):181-191(2008)
Chen, G., et al. Cancer Res. 67(19):9024-9029(2007)
Kaivosaari, S., et al. Mol. Pharmacol. 72(3):761-768(2007)