

## **TBC1D1 Antibody (Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP17252c

### **Specification**

# TBC1D1 Antibody (Center) - Product Information

**Application** WB,E **Primary Accession 086TI0** Other Accession NP 055988.2 Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 133084 Antigen Region 617-645

### TBC1D1 Antibody (Center) - Additional Information

**Gene ID 23216** 

#### **Other Names**

TBC1 domain family member 1, TBC1D1, KIAA1108

#### Target/Specificity

This TBC1D1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 617-645 amino acids from the Central region of human TBC1D1.

# **Dilution**

WB~~1:1000

## **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

# **Precautions**

TBC1D1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

# TBC1D1 Antibody (Center) - Protein Information

Name TBC1D1

Synonyms KIAA1108



**Function** May act as a GTPase-activating protein for Rab family protein(s). May play a role in the cell cycle and differentiation of various tissues. Involved in the trafficking and translocation of GLUT4-containing vesicles and insulin-stimulated glucose uptake into cells (By similarity).


**Cellular Location** Nucleus.

# **TBC1D1** Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# TBC1D1 Antibody (Center) - Images




TBC1D1 Antibody (Center) (Cat. #AP17252c) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the TBC1D1 antibody detected the TBC1D1 protein (arrow).

### TBC1D1 Antibody (Center) - Background

TBC1D1 is the founding member of a family of proteins sharing a 180- to 200-amino acid TBC domain presumed to have a role in regulating cell growth and differentiation. These proteins share significant homology with TRE2 (USP6; MIM 604334), yeast Bub2, and CDC16 (MIM 603461) (White et al., 2000 [PubMed 10965142]).[supplied by OMIM].

### **TBC1D1** Antibody (Center) - References

Bailey, S.D., et al. Diabetes Care (2010) In press: Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010): Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009) Ishibashi, K., et al. Genes Cells 14(1):41-52(2009)





Must, A., et al. Neuromolecular Med. 11(1):13-19(2009)