

AP4M1 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP17446c

Specification

AP4M1 Antibody (Center) - Product Information

Application WB,E
Primary Accession 000189

Other Accession Q2PWT8, Q9IKC7, Q29RY8, NP 004713.2

Reactivity Human

Predicted Bovine, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 49977
Antigen Region 223-251

AP4M1 Antibody (Center) - Additional Information

Gene ID 9179

Other Names

AP-4 complex subunit mu-1, AP-4 adaptor complex mu subunit, Adaptor-related protein complex 4 subunit mu-1, Mu subunit of AP-4, Mu-adaptin-related protein 2, mu-ARP2, Mu4-adaptin, mu4, AP4M1, MUARP2

Target/Specificity

This AP4M1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 223-251 amino acids from the Central region of human AP4M1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

AP4M1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

AP4M1 Antibody (Center) - Protein Information

Name AP4M1 (HGNC:574)



Synonyms MUARP2

Function Component of the adaptor protein complex 4 (AP-4). Adaptor protein complexes are vesicle coat components involved both in vesicle formation and cargo selection. They control the vesicular transport of proteins in different trafficking pathways (PubMed:10436028, PubMed:11139587, PubMed:10066790, PubMed:11802162, PubMed:20230749). AP-4 forms a non clathrin-associated coat on vesicles departing the trans-Golgi network (TGN) and may be involved in the targeting of proteins from the trans-Golgi network (TGN) to the endosomal-lysosomal system (PubMed:11139587, PubMed:20230749). It is also involved in protein sorting to the basolateral membrane in epithelial cells and the proper asymmetric localization of somatodendritic proteins in neurons (By similarity). Within AP-4, the mu-type subunit AP4M1 is directly involved in the recognition and binding of tyrosine-based sorting signals found in the cytoplasmic part of cargos (PubMed:10436028, PubMed:11139587, PubMed:26544806, PubMed:20230749). The adaptor protein complex 4 (AP-4) may also recognize other types of sorting signal (By similarity).

Cellular Location

Golgi apparatus, trans-Golgi network membrane; Peripheral membrane protein. Early endosome. Note=Found in soma and dendritic shafts of neuronal cells. {ECO:0000250|UniProtKB:Q2PWT8}

Tissue Location


Ubiquitous. Highly expressed in testis and lowly expressed in brain and lung.

AP4M1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

AP4M1 Antibody (Center) - Images



AP4M1 Antibody (Center) (Cat. #AP17446c) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the AP4M1 antibody detected the AP4M1 protein (arrow).



AP4M1 Antibody (Center) - Background

This gene encodes a subunit of the heterotetrameric AP-4 complex. The encoded protein belongs to the adaptor complexes medium subunits family. This AP-4 complex is involved in the recognition and sorting of cargo proteins with tyrosine-based motifs from the trans-golgi network to the endosomal-lysosomal system.

AP4M1 Antibody (Center) - References

Verkerk, A.J., et al. Am. J. Hum. Genet. 85(1):40-52(2009) Matsuda, S., et al. Neuron 57(5):730-745(2008) Lamesch, P., et al. Genomics 89(3):307-315(2007) Simmen, T., et al. Nat. Cell Biol. 4(2):154-159(2002) Boehm, M., et al. EMBO J. 20(22):6265-6276(2001)