

MRPS2 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP17471a

Specification

MRPS2 Antibody (N-term) - Product Information

| Application | WB,E |
|-------------------|--------------------|
| Primary Accession | <u>Q9Y399</u> |
| Other Accession | <u>NP_057118.1</u> |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Calculated MW | 33249 |
| Antigen Region | 44-71 |
| | |

MRPS2 Antibody (N-term) - Additional Information

Gene ID 51116

Other Names 28S ribosomal protein S2, mitochondrial, MRP-S2, S2mt, MRPS2

Target/Specificity This MRPS2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 44-71 amino acids from the N-terminal region of human MRPS2.

Dilution WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions MRPS2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MRPS2 Antibody (N-term) - Protein Information

Name MRPS2

Function Required for mitoribosome formation and stability, and mitochondrial translation.




Cellular Location Mitochondrion.

MRPS2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

MRPS2 Antibody (N-term) - Images



MRPS2 Antibody (N-term) (Cat. #AP17471a) western blot analysis in HepG2 cell line lysates (35ug/lane).This demonstrates the MRPS2 antibody detected the MRPS2 protein (arrow).

MRPS2 Antibody (N-term) - Background

Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein that belongs to the ribosomal protein S2 family.

MRPS2 Antibody (N-term) - References

Zhang, Z., et al. Genomics 81(5):468-480(2003) Kenmochi, N., et al. Genomics 77 (1-2), 65-70 (2001) :



Suzuki, T., et al. J. Biol. Chem. 276(35):33181-33195(2001) Cavdar Koc, E., et al. J. Biol. Chem. 276(22):19363-19374(2001)