10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

RAB32 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17478a

Specification

RAB32 Antibody (N-term) - Product Information

Application	**WB,E**
Primary Accession	Q13637
Other Accession	NP_006825.1
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**24997**
Antigen Region	**2-28**

RAB32 Antibody (N-term) - Additional Information

Gene ID 10981

Other Names
Ras-related protein Rab-32, RAB32

Target/Specificity
This RAB32 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 2-28 amino acids from the N-terminal region of human RAB32.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
RAB32 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

RAB32 Antibody (N-term) - Protein Information

Name RAB32

Function Acts as an A-kinase anchoring protein by binding to the type II regulatory subunit of protein kinase A and anchoring it to the mitochondrion. Also involved in synchronization of

mitochondrial fission (PubMed:12186851). Plays a role in the maturation of phagosomes that engulf pathogens, such as S.aureus and M.tuberculosis (PubMed:21255211). Plays an important role in the control of melanin production and melanosome biogenesis (PubMed:23084991). In concert with RAB38, regulates the proper trafficking of melanogenic enzymes TYR, TYRP1 and DCT/TYRP2 to melanosomes in melanocytes (By similarity).

Cellular Location
Mitochondrion. Mitochondrion outer membrane; Lipid-anchor. Cytoplasmic vesicle, phagosome. Cytoplasmic vesicle, phagosome membrane; Lipid-anchor; Cytoplasmic side. Melanosome {ECO:0000250|UniProtKB:Q9CZE3}. Melanosome membrane. Note=Recruited to phagosomes containing S.aureus or M.tuberculosis (PubMed:21255211). The BLOC-3 complex, a heterodimer of HPS1 and HPS4 promotes its membrane localization (PubMed:23084991).


Tissue Location
Widely expressed with high levels in heart, liver, kidney, bone marrow, testis, colon and fetal lung

RAB32 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

RAB32 Antibody (N-term) - Images



Anti-RAB32 Antibody (N-term) at 1:1000 dilution + SK-BR-3 whole cell lysate. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 25 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

RAB32 Antibody (N-term) - Background

Small GTP-binding proteins of the RAB family, such as RAB32, play essential roles in vesicle and granule targeting (Bao et al., 2002 [PubMed 11784320]).

RAB32 Antibody (N-term) - References

Hirota, Y., et al. Cell. Mol. Life Sci. 66(17):2913-2932(2009)
Shibata, D., et al. Int. J. Cancer 119(4):801-806(2006)
Mungall, A.J., et al. Nature 425(6960):805-811(2003)
Alto, N.M., et al. J. Cell Biol. 158(4):659-668(2002)
Bao, X., et al. Eur. J. Biochem. 269(1):259-271(2002)