

AIDA Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17627b**Specification**

AIDA Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q96BJ3
Other Accession	Q8C4Q6 , Q4R8C7 , Q6PBN2 , NP_073742.2
Reactivity	Human
Predicted	Zebrafish, Monkey, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35023
Antigen Region	255-283

AIDA Antibody (C-term) - Additional Information**Gene ID** 64853**Other Names**

Axin interactor, dorsalization-associated protein, Axin interaction partner and dorsalization antagonist, AIDA, C1orf80

Target/Specificity

This AIDA antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 255-283 amino acids from the C-terminal region of human AIDA.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

AIDA Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

AIDA Antibody (C-term) - Protein Information**Name** AIDA

Synonyms C1orf80

Function Acts as a ventralizing factor during embryogenesis. Inhibits axin-mediated JNK activation by binding axin and disrupting axin homodimerization. This in turn antagonizes a Wnt/beta-catenin- independent dorsalization pathway activated by AXIN/JNK-signaling (By similarity).


Tissue Location

Widely expressed in adult tissues, with highest expression in the heart and skeletal muscle

AIDA Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

AIDA Antibody (C-term) - Images

AIDA Antibody (C-term) (Cat. #AP17627b) western blot analysis in MDA-MB453 cell line lysates (35ug/lane). This demonstrates the AIDA antibody detected the AIDA protein (arrow).

AIDA Antibody (C-term) - Background

AIDA acts as a ventralizing factor during embryogenesis. Inhibits axin-mediated JNK activation by binding axin and disrupting axin homodimerization. This in turn antagonizes a Wnt/beta-catenin-independent dorsalization pathway activated by AXIN/JNK-signaling (By similarity).

AIDA Antibody (C-term) - References

Rui, Y., et al. Dev. Cell 13(2):268-282(2007)