

OR2Y1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP17842b

Specification

OR2Y1 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	<u>Q8NGV0</u>
Other Accession	<u>NP_001001657.1</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	34731
Antigen Region	244-271

OR2Y1 Antibody (C-term) - Additional Information

Gene ID 134083

Other Names Olfactory receptor 2Y1, Olfactory receptor OR5-2, OR2Y1

Target/Specificity This OR2Y1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 244-271 amino acids from the C-terminal region of human OR2Y1.

Dilution WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions OR2Y1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OR2Y1 Antibody (C-term) - Protein Information

Name OR2Y1

Function Odorant receptor.



Cellular Location


Cell membrane; Multi-pass membrane protein.

OR2Y1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

OR2Y1 Antibody (C-term) - Images



OR2Y1 Antibody (C-term) (Cat. #AP17842b) western blot analysis in NCI-H460 cell line lysates (35ug/lane).This demonstrates the OR2Y1 antibody detected the OR2Y1 protein (arrow).

OR2Y1 Antibody (C-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR2Y1 Antibody (C-term) - References

Johansson, A., et al. Obesity (Silver Spring) 18(4):803-808(2010) Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004) Fuchs, T., et al. Genomics 80(3):295-302(2002)