

ACCN5 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP17951b

Specification

ACCN5 Antibody (C-term) - Product Information

Application WB,E **Primary Accession O9NY37** NP 059115.1 Other Accession Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 57464 Antigen Region 388-414

ACCN5 Antibody (C-term) - Additional Information

Gene ID 51802

Other Names

Acid-sensing ion channel 5, ASIC5, Amiloride-sensitive cation channel 5, Human intestine Na(+) channel, HINaC, ASIC5, ACCN5, HINAC

Target/Specificity

This ACCN5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 388-414 amino acids from the C-terminal region of human ACCN5.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ACCN5 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ACCN5 Antibody (C-term) - Protein Information

Name ASIC5

Synonyms ACCN5, HINAC



Function Cation channel that gives rise to very low constitutive currents in the absence of activation. The activated channel exhibits selectivity for sodium, and is inhibited by amiloride.

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location


Detected in small intestine, duodenum and jejunum. Detected at very low levels in testis and rectum

ACCN5 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ACCN5 Antibody (C-term) - Images



ACCN5 Antibody (C-term) (Cat. #AP17951b) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the ACCN5 antibody detected the ACCN5 protein (arrow).

ACCN5 Antibody (C-term) - Background

This gene belongs to the amiloride-sensitive Na+ channel and degenerin (NaC/DEG) family, members of which have been identified in many animal species ranging from the nematode to human. The amiloride-sensitive Na(+) channel encoded by this gene is primarily expressed in the small intestine, however, its exact function is not known.

ACCN5 Antibody (C-term) - References

Schaefer, L., et al. FEBS Lett. 471 (2-3), 205-210 (2000) :