

## **GPR20 Antibody (C-term)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP17975b

### **Specification**

# **GPR20 Antibody (C-term) - Product Information**

WB,E Application **Primary Accession** 099678 Other Accession NP 005284.2 Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 38650 Antigen Region 298-324

### GPR20 Antibody (C-term) - Additional Information

#### **Gene ID 2843**

### **Other Names**

G-protein coupled receptor 20, GPR20

### Target/Specificity

This GPR20 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 298-324 amino acids from the C-terminal region of human GPR20.

## **Dilution**

WB~~1:1000

## **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

# **Precautions**

GPR20 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## **GPR20 Antibody (C-term) - Protein Information**

## Name GPR20

Function Orphan receptor with constitutive G(i) signaling activity that activate cyclic AMP.



Tel: 858.875.1900 Fax: 858.875.1999

### **Cellular Location**

Cell membrane; Multi-pass membrane protein

#### **Tissue Location**


Ubiquitous with highest levels in intestinal tissues. In the brain detected in thalamus, putamen, and caudate, but not in frontal cortex, pons and hypothalamus

## **GPR20 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## GPR20 Antibody (C-term) - Images



GPR20 Antibody (C-term) (Cat. #AP17975b) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the GPR20 antibody detected the GPR20 protein (arrow).

### GPR20 Antibody (C-term) - Background

Orphan receptor with constitutive G(i) signaling activity that activate cyclic AMP.

## **GPR20 Antibody (C-term) - References**

Hase, M., et al. J. Biol. Chem. 283(19):12747-12755(2008) O'Dowd, B.F., et al. Gene 187(1):75-81(1997)