

BCL7C Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18065a**Specification**

BCL7C Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	O8WUZ0
Other Accession	O08664 , NP_004756.2
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	23468
Antigen Region	8-34

BCL7C Antibody (N-term) - Additional Information**Gene ID** 9274**Other Names**

B-cell CLL/lymphoma 7 protein family member C, BCL7C

Target/Specificity

This BCL7C antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 8-34 amino acids from the N-terminal region of human BCL7C.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BCL7C Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

BCL7C Antibody (N-term) - Protein Information**Name** BCL7C**Function** May play an anti-apoptotic role.


Tissue Location

Ubiquitous..

BCL7C Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BCL7C Antibody (N-term) - Images

BCL7C Antibody (N-term) (Cat. #AP18065a) western blot analysis in SK-BR-3 cell line lysates (35ug/lane). This demonstrates the BCL7C antibody detected the BCL7C protein (arrow).

BCL7C Antibody (N-term) - Background

This gene is identified by the similarity of its product to the N-terminal region of BCL7A protein. The BCL7A protein is encoded by the gene known to be directly involved in a three-way gene translocation in a Burkitt lymphoma cell line. The function of this gene has not yet been determined.

BCL7C Antibody (N-term) - References

- Hosgood, H.D. III, et al. Occup Environ Med 66(12):848-853(2009)
Liang, X.S., et al. Br. J. Haematol. 146(4):418-423(2009)
Morton, L.M., et al. Cancer Epidemiol. Biomarkers Prev. 18(4):1259-1270(2009)
Lamesch, P., et al. Genomics 89(3):307-315(2007)
Olsen, J.V., et al. Cell 127(3):635-648(2006)