

DUS2L Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18265b

Specification

DUS2L Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O9NX74
Other Accession	NP_060273.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	55050
Antigen Region	371-397

DUS2L Antibody (C-term) - Additional Information

Gene ID 54920

Other Names

tRNA-dihydrouridine(20) synthase [NAD(P)+]-like, 131-, Dihydrouridine synthase 2, Up-regulated in lung cancer protein 8, URLC8, tRNA-dihydrouridine synthase 2-like, hDUS2, DUS2, DUS2L

Target/Specificity

This DUS2L antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 371-397 amino acids from the C-terminal region of human DUS2L.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DUS2L Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

DUS2L Antibody (C-term) - Protein Information

Name DUS2

Synonyms DUS2L

Function Dihydrouridine synthase. Catalyzes the NADPH-dependent synthesis of dihydrouridine, a modified base found in the D-loop of most tRNAs (PubMed:[15994936](#), PubMed:[26429968](#), PubMed:[30149704](#), PubMed:[34798057](#)). Negatively regulates the activation of EIF2AK2/PKR (PubMed:[18096616](#)).

Cellular Location

Cytoplasm. Endoplasmic reticulum. Note=Mainly at the endoplasmic reticulum.

Tissue Location


Weak expression in heart, placenta and skeletal muscle. Up-regulated in most lung cancer cells (at protein level)

DUS2L Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DUS2L Antibody (C-term) - Images



DUS2L Antibody (C-term) (Cat. #AP18265b) western blot analysis in A549 cell line lysates (35ug/lane). This demonstrates the DUS2L antibody detected the DUS2L protein (arrow).

DUS2L Antibody (C-term) - Background

Dihydrouridine synthase catalyzes reduction of the 5,6-double bond of a uridine residue on the displacement loop of tRNA. The resultant modified base, 5,6-dihydrouridine, appears to increase the conformational flexibility and dynamic motion of tRNA (Kato et al., 2005 [PubMed 15994936]).

DUS2L Antibody (C-term) - References

Mittelstadt, M., et al. Nucleic Acids Res. 36(3):998-1008(2008)
Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)
Lamesch, P., et al. Genomics 89(3):307-315(2007)
Kato, T., et al. Cancer Res. 65(13):5638-5646(2005)