

UVRAG Antibody (C-term L555)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1850e**Specification**

UVRAG Antibody (C-term L555) - Product Information

Application	WB,E
Primary Accession	Q9P2Y5
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	540-569

UVRAG Antibody (C-term L555) - Additional Information**Gene ID** 7405**Other Names**

UV radiation resistance-associated gene protein, p63, UVRAG

Target/Specificity

This UVRAG antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 540-569 amino acids from the C-terminal region of human UVRAG.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

UVRAG Antibody (C-term L555) is for research use only and not for use in diagnostic or therapeutic procedures.

UVRAG Antibody (C-term L555) - Protein Information**Name** UVRAG

Function Versatile protein that is involved in regulation of different cellular pathways implicated in membrane trafficking. Involved in regulation of the COPI-dependent retrograde transport from Golgi and the endoplasmic reticulum by associating with the NRZ complex; the function is dependent on its binding to phosphatidylinositol 3- phosphate (PtdIns(3)P) (PubMed:[16799551](#),

PubMed:[18552835](#), PubMed:[20643123](#), PubMed:[24056303](#), PubMed:[28306502](#)). During autophagy acts as a regulatory subunit of the alternative PI3K complex II (PI3KC3-C2) that mediates formation of phosphatidylinositol 3-phosphate and is believed to be involved in maturation of autophagosomes and endocytosis. Activates lipid kinase activity of PIK3C3 (PubMed:[16799551](#), PubMed:[20643123](#), PubMed:[24056303](#), PubMed:[28306502](#)). Involved in the regulation of degradative endocytic trafficking and cytokinesis, and in regulation of ATG9A transport from the Golgi to the autophagosome; the functions seems to implicate its association with PI3KC3-C2 (PubMed:[16799551](#), PubMed:[20643123](#), PubMed:[24056303](#)). Involved in maturation of autophagosomes and degradative endocytic trafficking independently of BECN1 but depending on its association with a class C Vps complex (possibly the HOPS complex); the association is also proposed to promote autophagosome recruitment and activation of Rab7 and endosome-endosome fusion events (PubMed:[18552835](#), PubMed:[28306502](#)). Enhances class C Vps complex (possibly HOPS complex) association with a SNARE complex and promotes fusogenic SNARE complex formation during late endocytic membrane fusion (PubMed:[24550300](#)). In case of negative- strand RNA virus infection is required for efficient virus entry, promotes endocytic transport of virions and is implicated in a VAMP8- specific fusogenic SNARE complex assembly (PubMed:[24550300](#)).

Cellular Location

Late endosome. Lysosome. Cytoplasmic vesicle, autophagosome. Early endosome. Endoplasmic reticulum. Midbody. Chromosome, centromere. Note=Colocalizes with RAB9-positive compartments involved in retrograde transport from late endosomes to trans-Golgi network. Colocalization with early endosomes is only partial (PubMed:[24056303](#)). Recruited to autophagosome following interaction with RUBCNL/PACER (PubMed:[28306502](#))

Tissue Location


Highly expressed in brain, lung, kidney and liver.


UVRAG Antibody (C-term L555) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UVRAG Antibody (C-term L555) - Images





Western blot analysis of UVRAG Antibody (C-term L555) (Cat.#AP1850e) in Ramos cell line lysates (35ug/lane). UVRAG (arrow) was detected using the purified Pab.

UVRAG Antibody (C-term L555) - Background

UVRAG complements the ultraviolet sensitivity of xeroderma pigmentosum group C cells and encodes a protein with a C2 domain. The protein activates the Beclin1-PI(3)KC3 complex, promoting autophagy and suppressing the proliferation and tumorigenicity of human colon cancer cells. Chromosomal aberrations involving the UVAG gene are associated with left-right axis malformation and mutations in the gene have been associated with colon cancer.

UVRAG Antibody (C-term L555) - References

Liang,C., et al. Nat. Cell Biol. 8 (7), 688-699 (2006)
Ionov,Y., et al. Oncogene 23 (3), 639-645 (2004)
Goi,T., et al. Surg. Today 33 (9), 702-706 (2003)
Iida,A., et al. Hum. Genet. 106 (3), 277-287 (2000)
Perelman,B., et al. Genomics 41 (3), 397-405 (1997)
Bekri,S., et al. Cytogenet. Cell Genet. 79 (1-2), 125-131 (1997)
Teitz,T., et al. Gene 87 (2), 295-298 (1990)

UVRAG Antibody (C-term L555) - Citations

- [Autophagic induction of amyotrophic lateral sclerosis-linked Cu/Zn superoxide dismutase 1 G93A mutant in NSC34 cells.](#)
- [RalB and the exocyst mediate the cellular starvation response by direct activation of autophagosome assembly.](#)