

IL17RC Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18560a

Specification

IL17RC Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q8NAC3
Other Accession	A1L1A6 , NP_116121.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	86240
Antigen Region	103-131

IL17RC Antibody (N-term) - Additional Information

Gene ID 84818

Other Names

Interleukin-17 receptor C, IL-17 receptor C, IL-17RC, Interleukin-17 receptor homolog, IL17Rhom, Interleukin-17 receptor-like protein, IL-17RL, ZcytoR14, IL17RC

Target/Specificity

This IL17RC antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 103-131 amino acids from the N-terminal region of human IL17RC.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IL17RC Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

IL17RC Antibody (N-term) - Protein Information

Name IL17RC

Function Receptor for IL17A and IL17F, major effector cytokines of innate and adaptive immune

system involved in antimicrobial host defense and maintenance of tissue integrity (By similarity). Receptor for IL17A and IL17F, major effector cytokines of innate and adaptive immune system involved in antimicrobial host defense and maintenance of tissue integrity. Receptor for IL17A and IL17F homodimers as part of a heterodimeric complex with IL17RA (PubMed:[16785495](#)). Receptor for the heterodimer formed by IL17A and IL17B as part of a heterodimeric complex with IL17RA (PubMed:[18684971](#)). Has also been shown to be the cognate receptor for IL17F and to bind IL17A with high affinity without the need for IL17RA (PubMed:[17911633](#)). Upon binding of IL17F homodimer triggers downstream activation of TRAF6 and NF-kappa-B signaling pathway (PubMed:[16785495](#), PubMed:[32187518](#)). Induces transcriptional activation of IL33, a potent cytokine that stimulates group 2 innate lymphoid cells and adaptive T-helper 2 cells involved in pulmonary allergic response to fungi (By similarity). Promotes sympathetic innervation of peripheral organs by coordinating the communication between gamma-delta T cells and parenchymal cells. Stimulates sympathetic innervation of thermogenic adipose tissue by driving TGFB1 expression (By similarity). Binding of IL17A-IL17F to IL17RA-IL17RC heterodimeric receptor complex triggers homotypic interaction of IL17RA and IL17RC chains with TRAF3IP2 adapter through SEFIR domains. This leads to downstream TRAF6-mediated activation of NF-kappa-B and MAPkinase pathways ultimately resulting in transcriptional activation of cytokines, chemokines, antimicrobial peptides and matrix metalloproteinases, with potential strong immune inflammation (PubMed:[18684971](#), PubMed:[17911633](#)). Primarily induces neutrophil activation and recruitment at infection and inflammatory sites (By similarity). Stimulates the production of antimicrobial beta-defensins DEFB1, DEFB103A, and DEFB104A by mucosal epithelial cells, limiting the entry of microbes through the epithelial barriers (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Note=Soluble isoforms may be produced

Tissue Location


Expressed in prostate, skeletal muscle, kidney and placenta (at protein level) (PubMed:11706037). Expressed in brain, cartilage, colon, heart, intestine, kidney, liver, lung, muscle, placenta, and prostate (PubMed:11706037). Also detected in thyroid, trachea and adrenal gland (PubMed:17911633). Low expression in thymus and leukocytes (PubMed:11706037).


IL17RC Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IL17RC Antibody (N-term) - Images





IL17RC Antibody (N-term) (Cat. #AP18560a) western blot analysis in Jurkat cell line lysates (35ug/lane). This demonstrates the IL17RC antibody detected the IL17RC protein (arrow).

IL17RC Antibody (N-term) - Background

This gene encodes a single-pass transmembrane protein that shares limited similarity with the interleukin-17 receptor. Multiple alternatively spliced transcript variants encoding different isoforms have been detected for this gene, but the full-length nature of only three have been determined to date.

IL17RC Antibody (N-term) - References

Pickens, S.R., et al. J. Immunol. 184(6):3233-3241(2010)
Wright, J.F., et al. J. Immunol. 181(4):2799-2805(2008)
Zrioual, S., et al. J. Immunol. 180(1):655-663(2008)
Kuestner, R.E., et al. J. Immunol. 179(8):5462-5473(2007)
Haudenschild, D.R., et al. Prostate 66(12):1268-1274(2006)