

PCDHGB1 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18632B

Specification

PCDHGB1 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O9Y5G3
Other Accession	NP_061745.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	100360
Antigen Region	773-799

PCDHGB1 Antibody (C-term) - Additional Information

Gene ID 56104

Other Names

Protocadherin gamma-B1, PCDH-gamma-B1, PCDHGB1

Target/Specificity

This PCDHGB1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 773-799 amino acids from the C-terminal region of human PCDHGB1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PCDHGB1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PCDHGB1 Antibody (C-term) - Protein Information

Name PCDHGB1

Function Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.


Cellular Location

Cell membrane; Single-pass type I membrane protein

PCDHGB1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PCDHGB1 Antibody (C-term) - Images

PCDHGB1 Antibody (C-term) (Cat. #AP18632b) western blot analysis in ZR-75-1 cell line lysates (35ug/lane). This demonstrates the PCDHGB1 antibody detected the PCDHGB1 protein (arrow).

PCDHGB1 Antibody (C-term) - Background

This gene is a member of the protocadherin gamma gene cluster, one of three related clusters tandemly linked on chromosome five. These gene clusters have an immunoglobulin-like organization, suggesting that a novel mechanism may be involved in their regulation and expression. The gamma gene cluster includes 22 genes divided into 3 subfamilies. Subfamily A contains 12 genes, subfamily B contains 7 genes and 2 pseudogenes, and the more distantly related subfamily C contains 3 genes. The tandem array of 22 large, variable region exons are followed by a constant region, containing 3 exons shared by all genes in the cluster. Each variable region exon encodes the extracellular region, which includes 6 cadherin ectodomains and a transmembrane region. The constant region exons encode the common cytoplasmic region. These neural cadherin-like cell adhesion proteins most likely play a critical role in the establishment and function of specific cell-cell connections in the brain. Alternative splicing has been described for the gamma cluster genes.

PCDHGB1 Antibody (C-term) - References

Wu, Q., et al. Genome Res. 11(3):389-404(2001)
Nollet, F., et al. J. Mol. Biol. 299(3):551-572(2000)
Yagi, T., et al. Genes Dev. 14(10):1169-1180(2000)
Wu, Q., et al. Proc. Natl. Acad. Sci. U.S.A. 97(7):3124-3129(2000)
Wu, Q., et al. Cell 97(6):779-790(1999)