10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

USP39 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18677c

Specification

USP39 Antibody (Center) - Product Information

Application	**WB,E**
Primary Accession	Q53GS9
Other Accession	Q3TIX9, NP_006581.2
Reactivity	**Human**
Predicted	**Mouse**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**65381**
Antigen Region	**237-265**

USP39 Antibody (Center) - Additional Information

Gene ID 10713

Other Names
U4/U6U5 tri-snRNP-associated protein 2, Inactive ubiquitin-specific peptidase 39, SAD1 homolog, U4/U6U5 tri-snRNP-associated 65 kDa protein, 65K, USP39

Target/Specificity
This USP39 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 237-265 amino acids from the Central region of human USP39.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
USP39 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

USP39 Antibody (Center) - Protein Information

Name USP39 (HGNC:20071)

Function Deubiquitinating enzyme that plays a role in many cellular processes including cellular antiviral response, epithelial morphogenesis, DNA repair or B-cell development (PubMed:33127822, PubMed:34614178). Plays a role in pre-mRNA splicing as a component of the U4/U6-U5 tri-snRNP, one of the building blocks of the precatalytic spliceosome (PubMed:11350945, PubMed:26912367). Specifically regulates immunoglobulin gene rearrangement in a spliceosome-dependent manner, which involves modulating chromatin interactions at the Igh locus and therefore plays an essential role in B-cell development (By similarity). Regulates AURKB mRNA levels, and thereby plays a role in cytokinesis and in the spindle checkpoint (PubMed:18728397). Regulates apoptosis and G2/M cell cycle checkpoint in response to DNA damage by deubiquitinating and stabilizing CHK2 (PubMed:30771428). Plays also an important role in DNA repair by controlling the recruitment of XRCC4/LIG4 to DNA double-strand breaks for non-homologous end-joining repair (PubMed:34614178). Participates in antiviral activity by affecting the type I IFN signaling by stabilizing STAT1 and decreasing its 'Lys-6'-linked ubiquitination (PubMed:33127822). Contributes to non-canonical Wnt signaling during epidermal differentiation (By similarity). Acts as a negative regulator NF-kappa-B activation through deubiquitination of 'Lys-48'-linked ubiquitination of NFKBIA (PubMed:36651806).


Cellular Location
Nucleus

USP39 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

USP39 Antibody (Center) - Images



USP39 Antibody (Center) (Cat. #AP18677c) western blot analysis in Hela cell line lysates (35ug/lane).This demonstrates the USP39 antibody detected the USP39 protein (arrow).

USP39 Antibody (Center) - Background

USP39 may play a role in mRNA splicing. It is unsure if the protein really exhibits hydrolase activity. Could be a competitor of ubiquitin C-terminal hydrolases (UCHs).

USP39 Antibody (Center) - References

Rose, J. Phd, et al. Mol. Med. (2010) In press :
van Leuken, R.J., et al. Cell Cycle 7(17):2710-2719(2008)
Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)
Ewing, R.M., et al. Mol. Syst. Biol. 3, 89 (2007) :
Olsen, J.V., et al. Cell 127(3):635-648(2006)