

**Mouse Il17ra Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP18687b**

**Specification**

---

**Mouse Il17ra Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q60943</a>
Other Accession	<a href="#">NP_032385.1</a>
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	721-747

**Mouse Il17ra Antibody (C-term) - Additional Information**

**Gene ID** 16172

**Other Names**

Interleukin-17 receptor A, IL-17 receptor A, IL-17RA, CD217, Il17ra, Il17r

**Target/Specificity**

This Mouse Il17ra antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 721-747 amino acids of mouse Il17ra.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Mouse Il17ra Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**Mouse Il17ra Antibody (C-term) - Protein Information**

**Name** Il17ra

**Synonyms** Il17r

**Function** Receptor for IL17A and IL17F, major effector cytokines of innate and adaptive immune

system involved in antimicrobial host defense and maintenance of tissue integrity. Receptor for IL17A (PubMed:[17911633](#), PubMed:[20554964](#), PubMed:[8777726](#), PubMed:[27923703](#)). Receptor for IL17F (PubMed:[17911633](#), PubMed:[20554964](#)). Binds to IL17A with higher affinity than to IL17F (PubMed:[17911633](#)). Binds IL17A and IL17F homodimers as part of a heterodimeric complex with IL17RC (By similarity). Also binds heterodimers formed by IL17A and IL17F as part of a heterodimeric complex with IL17RC (By similarity). Cytokine binding triggers homotypic interaction of IL17RA and IL17RC chains with TRAF3IP2 adapter, leading to TRAF6-mediated activation of NF-kappa-B and MAPkinase pathways, ultimately resulting in transcriptional activation of cytokines, chemokines, antimicrobial peptides and matrix metalloproteinases, with potential strong immune inflammation (By similarity). Involved in antimicrobial host defense primarily promoting neutrophil activation and recruitment at infection sites to destroy extracellular bacteria and fungi (PubMed:[21993848](#), PubMed:[20364087](#)). In secondary lymphoid organs, contributes to germinal center formation by regulating the chemotactic response of B cells to CXCL12 and CXCL13, enhancing retention of B cells within the germinal centers, B cell somatic hypermutation rate and selection toward plasma cells (PubMed:[18157131](#)). Plays a role in the maintenance of the integrity of epithelial barriers during homeostasis and pathogen infection. Stimulates the production of antimicrobial beta-defensins DEFB1, DEFB103A, and DEFB104A by mucosal epithelial cells, limiting the entry of microbes through the epithelial barriers (PubMed:[19144317](#)). Involved in antiviral host defense through various mechanisms. Enhances immunity against West Nile virus by promoting T cell cytotoxicity (PubMed:[27795421](#)). Contributes to influenza A virus (H1N1) clearance by driving the differentiation of B-1a B cells, providing for production of virus-specific IgM antibodies at first line of host defense (PubMed:[26735852](#)). Receptor for IL17C as part of a heterodimeric complex with IL17RE (PubMed:[21993848](#), PubMed:[21993849](#), PubMed:[21982598](#)).

#### Cellular Location

Cell membrane; Single-pass type I membrane protein

#### Tissue Location


Widely expressed (PubMed:[21993848](#)). Highly expressed in T cells and macrophages (PubMed:[19144317](#)). Highly expressed in B-1a B cells and at a lower extent in B-1b and B-2 B cells (at protein level) (PubMed:[26735852](#)).


#### Mouse IL17ra Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Mouse IL17ra Antibody (C-term) - Images





Mouse Il17ra Antibody (C-term) (Cat. #AP18687b) western blot analysis in mouse NIH-3T3 cell line lysates (35ug/lane). This demonstrates the Mouse Il17ra antibody detected the Mouse Il17ra protein (arrow).

#### **Mouse Il17ra Antibody (C-term) - Background**

Receptor for IL17A. Binds its ligand with low affinity, suggesting that additional components are involved in IL17A-induced signaling (By similarity).

#### **Mouse Il17ra Antibody (C-term) - References**

Onishi, R.M., et al. J. Biol. Chem. 285(43):32751-32759(2010)  
Guiton, R., et al. J. Infect. Dis. 202(3):427-435(2010)  
Mitsdoerffer, M., et al. Proc. Natl. Acad. Sci. U.S.A. 107(32):14292-14297(2010)  
Hill, G.R., et al. Blood 116(5):819-828(2010)  
Wu, H.J., et al. Immunity 32(6):815-827(2010)