

GSC Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP18734a

Specification

GSC Antibody (N-term) - Product Information

Application WB,E
Primary Accession P56915

Other Accession <u>Q02591</u>, <u>NP 776248.1</u>

Reactivity
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Mouse
Rabbit
Polyclonal
Rabbit IgG
28150
38-65

GSC Antibody (N-term) - Additional Information

Gene ID 145258

Other Names

Homeobox protein goosecoid, GSC

Target/Specificity

This GSC antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 38-65 amino acids from the N-terminal region of human GSC.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GSC Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

GSC Antibody (N-term) - Protein Information

Name GSC

Function Regulates chordin (CHRD). May play a role in spatial programing within discrete embryonic fields or lineage compartments during organogenesis. In concert with NKX3-2, plays a



role in defining the structural components of the middle ear; required for the development of the entire tympanic ring (By similarity). Probably involved in the regulatory networks that define neural crest cell fate specification and determine mesoderm cell lineages in mammals.


Cellular Location Nucleus.

GSC Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

GSC Antibody (N-term) - Images



GSC Antibody (N-term)(Cat. #AP18734a) western blot analysis in mouse stomach tissue lysates (35ug/lane). This demonstrates the GSC antibody detected the GSC protein (arrow).

GSC Antibody (N-term) - Background

This gene encodes a member of the bicoid subfamily of the paired (PRD) homeobox family of proteins. The encoded protein acts as a transcription factor and may be autoregulatory. A similar protein in mice plays a role in craniofacial and rib cage development during embryogenesis.

GSC Antibody (N-term) - References

Zhang, Q., et al. Plast. Reconstr. Surg. 125(3):979-987(2010)
Paterson, A.D., et al. Diabetes 59(2):539-549(2010)
Zhang, Q.G., et al. Plast. Reconstr. Surg. 124(4):1157-1164(2009)
Schlade-Bartusiak, K., et al. Am. J. Med. Genet. A 146A (1), 117-123 (2008):
Hartwell, K.A., et al. Proc. Natl. Acad. Sci. U.S.A. 103(50):18969-18974(2006)