

RRAGB Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18936A**Specification**

RRAGB Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q5VZM2
Other Accession	NP_006055.3
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	43250
Antigen Region	1-30

RRAGB Antibody (N-term) - Additional Information**Gene ID** 10325**Other Names**

Ras-related GTP-binding protein B, Rag B, RagB, RRAGB (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=19901)
target="_blank">HGNC:19901)

Target/Specificity

This RRAGB antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human RRAGB.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RRAGB Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

RRAGB Antibody (N-term) - Protein Information**Name** RRAGB ([HGNC:19901](#))

Function Guanine nucleotide-binding protein that plays a crucial role in the cellular response to amino acid availability through regulation of the mTORC1 signaling cascade (PubMed:[18497260](#), PubMed:[20381137](#), PubMed:[23723238](#), PubMed:[24095279](#)). Forms heterodimeric Rag complexes with RagC/RRAGC or RagD/RRAGD and cycles between an inactive GDP-bound and an active GTP-bound form: RagB/RRAGB is in its active form when GTP-bound RagB/RRAGB forms a complex with GDP-bound RagC/RRAGC (or RagD/RRAGD) and in an inactive form when GDP-bound RagB/RRAGB heterodimerizes with GTP-bound RagC/RRAGC (or RagD/RRAGD) (PubMed:[18497260](#), PubMed:[20381137](#), PubMed:[23723238](#), PubMed:[24095279](#)). In its GTP-bound active form, promotes the recruitment of mTORC1 to the lysosomes and its subsequent activation by the GTPase RHEB (PubMed:[18497260](#), PubMed:[20381137](#), PubMed:[23723238](#)). Involved in the RCC1/Ran-GTPase pathway (PubMed:[9394008](#)).

Cellular Location


Cytoplasm. Lysosome membrane. Note=Recruited to the lysosome surface by the Ragulator complex.

RRAGB Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RRAGB Antibody (N-term) - Images



RRAGB Antibody (N-term) (Cat. #AP18936a) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the RRAGB antibody detected the RRAGB protein (arrow).

RRAGB Antibody (N-term) - Background

Ras-homologous GTPases constitute a large family of signal transducers that alternate between an activated, GTP-binding state and an inactivated, GDP-binding state. These proteins represent cellular switches that are operated by GTP-exchange factors and

factors that stimulate their intrinsic GTPase activity. All GTPases of the Ras superfamily have in common the presence of six conserved motifs involved in GTP/GDP binding, three of which are phosphate-/magnesium-binding sites (PM1-PM3) and three of which are guanine nucleotide-binding sites (G1-G3). Transcript variants encoding distinct isoforms have been identified. [provided by RefSeq].

RRAGB Antibody (N-term) - References

Ross, M.T., et al. Nature 434(7031):325-337(2005)
Tomarev, S.I., et al. Invest. Ophthalmol. Vis. Sci. 44(6):2588-2596(2003)
Sekiguchi, T., et al. J. Biol. Chem. 276(10):7246-7257(2001)
Hirose, E., et al. J. Cell. Sci. 111 (PT 1), 11-21 (1998) :
Schurmann, A., et al. J. Biol. Chem. 270(48):28982-28988(1995)