

KCNG2 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP18984c

Specification

KCNG2 Antibody (Center) - Product Information

Application WB,E **Primary Accession 09UI96** Other Accession NP 036415.1 Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 51240 Antigen Region 149-177

KCNG2 Antibody (Center) - Additional Information

Gene ID 26251

Other Names

Potassium voltage-gated channel subfamily G member 2, Cardiac potassium channel subunit, Voltage-gated potassium channel subunit Kv62, KCNG2, KCNF2

Target/Specificity

This KCNG2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 149-177 amino acids from the Central region of human KCNG2.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions


KCNG2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

KCNG2 Antibody (Center) - Protein Information

Name KCNG2

Synonyms KCNF2







Function Potassium channel subunit. Modulates channel activity by shifting the threshold and the half-maximal activation to more negative values.

Cellular Location

Membrane; Multi-pass membrane protein.

Tissue Location


Highly expressed in heart, liver, skeletal muscle, kidney and pancreas. Detected at low levels in brain, lung and placenta

KCNG2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

KCNG2 Antibody (Center) - Images



KCNG2 Antibody (Center) (Cat. #AP18984c) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the KCNG2 antibody detected the KCNG2 protein (arrow).

KCNG2 Antibody (Center) - Background

Voltage-gated potassium (Kv) channels represent the most complex class of voltage-gated ion channels from both functional and structural standpoints. Their diverse functions include regulating neurotransmitter release, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smooth muscle contraction, and cell volume. This gene encodes a member of the potassium channel, voltage-gated, subfamily G. This member is a gamma subunit of the voltage-gated potassium channel. The delayed-rectifier type channels containing this subunit may contribute to cardiac action potential repolarization. [provided by





Tel: 858.875.1900 Fax: 858.875.1999

RefSeq].

KCNG2 Antibody (Center) - References

Gutman, G.A., et al. Pharmacol. Rev. 57(4):473-508(2005) Zhu, X.R., et al. Recept. Channels 6(5):337-350(1999)