

PREP Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18992C

Specification

PREP Antibody (Center) - Product Information

Application	WB,E
Primary Accession	P48147
Other Accession	NP_002717.3
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	80700
Antigen Region	293-322

PREP Antibody (Center) - Additional Information

Gene ID 5550

Other Names

Prolyl endopeptidase, PE, Post-proline cleaving enzyme, PREP, PEP

Target/Specificity

This PREP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 293-322 amino acids from the Central region of human PREP.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PREP Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

PREP Antibody (Center) - Protein Information

Name PREP

Synonyms PEP

Function Cleaves peptide bonds on the C-terminal side of prolyl residues within peptides that are up to approximately 30 amino acids long.

Cellular Location


Cytoplasm.

PREP Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PREP Antibody (Center) - Images



PREP Antibody (Center) (Cat. #AP18992c) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the PREP antibody detected the PREP protein (arrow).

PREP Antibody (Center) - Background

The protein encoded by this gene is a cytosolic prolyl endopeptidase that cleaves peptide bonds on the C-terminal side of prolyl residues within peptides that are up to approximately 30 amino acids long. Prolyl endopeptidases have been reported to be involved in the maturation and degradation of peptide hormones and neuropeptides.

PREP Antibody (Center) - References

Pinheiro, A.P., et al. Am. J. Med. Genet. B Neuropsychiatr. Genet. 153B (5), 1070-1080 (2010) :
Rose, J. PhD, et al. Mol. Med. (2010) In press :
Tenorio-Laranga, J., et al. J Neuroinflammation 7, 23 (2010) :
Pillas, D., et al. PLoS Genet. 6 (2), E1000856 (2010) :
Malhotra, A., et al. Diabetes Metab. Res. Rev. 25(8):740-747(2009)