

PIG3 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19269b

Specification

PIG3 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q53FA7
Other Accession	NP_004872.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35536
Antigen Region	174-203

PIG3 Antibody (C-term) - Additional Information

Gene ID 9540

Other Names

Quinone oxidoreductase PIG3, 1---, Tumor protein p53-inducible protein 3, p53-induced gene 3 protein, TP53I3, PIG3

Target/Specificity

This PIG3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 174-203 amino acids from the C-terminal region of human PIG3.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PIG3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PIG3 Antibody (C-term) - Protein Information

Name TP53I3 ([HGNC:19373](#))

Synonyms PIG3


Function Catalyzes the NADPH-dependent reduction of quinones (PubMed:[19349281](#)). Exhibits a low enzymatic activity with beta- naphthoquinones, with a strong preference for the ortho-quinone isomer (1,2-beta-naphthoquinone) over the para isomer (1,4-beta- naphthoquinone). Also displays a low reductase activity for non-quinone compounds such as diamine and 2,6-dichloroindophenol (in vitro) (PubMed:[19349281](#)). Involved in the generation of reactive oxygen species (ROS) (PubMed:[19349281](#)).

PIG3 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PIG3 Antibody (C-term) - Images



PIG3 Antibody (C-term)(Cat. #AP19269b) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the PIG3 antibody detected the PIG3 protein (arrow).

PIG3 Antibody (C-term) - Background

The protein encoded by this gene is similar to oxidoreductases, which are enzymes involved in cellular responses to oxidative stresses and irradiation. This gene is induced by the tumor suppressor p53 and is thought to be involved in p53-mediated cell death. It contains a p53 consensus binding site in its promoter region and a downstream pentanucleotide microsatellite sequence. P53 has been shown to transcriptionally activate this gene by interacting with the downstream pentanucleotide microsatellite sequence. The microsatellite is polymorphic, with a varying number of pentanucleotide repeats directly correlated with the extent of transcriptional activation by p53. It has been

suggested that the microsatellite polymorphism may be associated with differential susceptibility to cancer. At least two transcript variants encoding the same protein have been found for this gene.

PIG3 Antibody (C-term) - References

Kotsinas, A., et al. Oncogene 29 (37), 5220 (2010) :
Liu, C.Y., et al. Carcinogenesis 31(7):1259-1263(2010)
Lee, J.H., et al. Oncogene 29(10):1431-1450(2010)
Guey, L.T., et al. Eur. Urol. 57(2):283-292(2010)
Hosgood, H.D. III, et al. Occup Environ Med 66(12):848-853(2009)