

WASH2P Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19270A

Specification

WASH2P Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q6VEQ5
Other Accession	C4AMC7 , B2RYF7 , Q8VDD8 , A7Z063
Reactivity	Human
Predicted	Bovine, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	50312
Antigen Region	1-30

WASH2P Antibody (N-term) - Additional Information

Other Names

WAS protein family homolog 2, CXYorf1-like protein on chromosome 2, Protein FAM39B, WASH2P, FAM39B

Target/Specificity

This WASH2P antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human WASH2P.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

WASH2P Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

WASH2P Antibody (N-term) - Protein Information

Name WASH2P

Synonyms FAM39B

Function Acts as a nucleation-promoting factor at the surface of endosomes, where it recruits and activates the Arp2/3 complex to induce actin polymerization, playing a key role in the fission of tubules that serve as transport intermediates during endosome sorting. Involved in endocytic trafficking of EGF. Involved in transferrin receptor recycling. Regulates the trafficking of endosomal alpha5beta1 integrin to the plasma membrane and involved in invasive cell migration. In T- cells involved in endosome-to-membrane recycling of receptors including T-cell receptor (TCR), CD28 and ITGAL; proposed to be implicated in T- cell proliferation and effector function. In dendritic cells involved in endosome-to-membrane recycling of major histocompatibility complex (MHC) class II probably involving retromer and subsequently allowing antigen sampling, loading and presentation during T-cell activation. Involved in Arp2/3 complex-dependent actin assembly driving Salmonella typhimurium invasion independent of ruffling. Involved in the exocytosis of MMP14 leading to matrix remodeling during invasive migration and implicating late endosome-to-plasma membrane tubular connections and cooperation with the exocyst complex. Involved in negative regulation of autophagy independently from its role in endosomal sorting by inhibiting BECN1 ubiquitination to inactivate PIK3C3/Vps34 activity (By similarity).

Cellular Location


Early endosome membrane {ECO:0000250|UniProtKB:A8K0Z3}. Recycling endosome membrane {ECO:0000250|UniProtKB:Q8VDD8}. Late endosome {ECO:0000250|UniProtKB:A8K0Z3}. Cytoplasmic vesicle, autophagosome {ECO:0000250|UniProtKB:Q8VDD8}. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole {ECO:0000250|UniProtKB:Q8VDD8}. Note=Localization to the endosome membrane is mediated via its interaction with WASHC2 {ECO:0000250|UniProtKB:A8K0Z3}

WASH2P Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

WASH2P Antibody (N-term) - Images



WASH2P Antibody (N-term)(Cat. #AP19270a) western blot analysis in MDA-MB453 cell line lysates (35ug/lane). This demonstrates the WASH2P antibody detected the WASH2P protein (arrow).

WASH2P Antibody (N-term) - Background

WASH2P acts as a nucleation-promoting factor at the surface of endosomes, where it recruits and activates the Arp2/3 complex to induce actin polymerization, playing a key role in the fission of tubules that serve as transport intermediates during endosome sorting (By similarity).