10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# MS4A13 Antibody(N-term)

**Affinity Purified Rabbit Polyclonal Antibody (Pab)**
**Catalog # AP19573a**

## Specification

### MS4A13 Antibody(N-term) - Product Information

| | |
|---|---|
| Application | **WB,E** |
| Primary Accession | Q5J8X5 |
| Other Accession | NP_001012417.2 |
| Reactivity | **Human, Mouse** |
| Host | **Rabbit** |
| Clonality | **Polyclonal** |
| Isotype | **Rabbit IgG** |
| Calculated MW | **17307** |
| Antigen Region | **7-36** |

### MS4A13 Antibody(N-term) - Additional Information

**Gene ID** 503497

**Other Names**
Membrane-spanning 4-domains subfamily A member 13, Testis-expressed transmembrane protein 4, MS4A13

**Target/Specificity**
This MS4A13 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 7-36 amino acids from the N-terminal region of human MS4A13.

**Dilution**
WB~~1:1000

**Format**
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**
MS4A13 Antibody(N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

### MS4A13 Antibody(N-term) - Protein Information

**Name** MS4A13

**Function** May be involved in signal transduction as a component of a multimeric receptor

complex.


**Cellular Location**
Membrane; Multi-pass membrane protein

## MS4A13 Antibody(N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## MS4A13 Antibody(N-term) - Images




MS4A13 Antibody (N-term) (Cat. #AP19573a) western blot analysis in mouse liver tissue lysates (35ug/lane).This demonstrates the MS4A13 antibody detected the MS4A13 protein (arrow).

## MS4A13 Antibody(N-term) - Background

MS4A13 may be involved in signal transduction as a component of a multimeric receptor complex (By similarity).

## MS4A13 Antibody(N-term) - References

Strausberg, R.L., et al. Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903(2002)