

OR52J3 Antibody(C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP19605b

Specification

OR52J3 Antibody(C-term) - Product Information

Application WB,E
Primary Accession Q8NH60

Other Accession NP 001001916.2

Reactivity
Human
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Human
Rabbit
Rabbit
Solution
Rabbit IgG
283-311

OR52J3 Antibody(C-term) - Additional Information

Gene ID 119679

Other Names

Olfactory receptor 52J3, Olfactory receptor OR11-32, OR52J3

Target/Specificity

This OR52J3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 283-311 amino acids from the C-terminal region of human OR52J3.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OR52J3 Antibody(C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OR52J3 Antibody(C-term) - Protein Information

Name OR52J3

Function Odorant receptor.



Cellular Location


Cell membrane; Multi-pass membrane protein.

OR52J3 Antibody(C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

OR52J3 Antibody(C-term) - Images




Anti-OR52J3 Antibody (C-term) at 1:1000 dilution + A2058 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 35 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

OR52J3 Antibody(C-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR52J3 Antibody(C-term) - References





Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)