

OR11A1 Antibody(C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19674b**Specification**

OR11A1 Antibody(C-term) - Product Information

Application	WB,E
Primary Accession	O9GZK7
Other Accession	NP_039225.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35250
Antigen Region	279-306

OR11A1 Antibody(C-term) - Additional Information**Gene ID** 26531**Other Names**

Olfactory receptor 11A1, Hs6M1-18, Olfactory receptor 11A2, Olfactory receptor OR6-30, OR11A1, OR11A2

Target/Specificity

This OR11A1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 279-306 amino acids from the C-terminal region of human OR11A1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OR11A1 Antibody(C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OR11A1 Antibody(C-term) - Protein Information**Name** OR11A1**Synonyms** OR11A2

Function Odorant receptor.

Cellular Location


Cell membrane; Multi-pass membrane protein.

OR11A1 Antibody(C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OR11A1 Antibody(C-term) - Images



OR11A1 Antibody (C-term) (Cat. #AP19674b) western blot analysis in MDA-MB453 cell line lysates (35ug/lane). This demonstrates the OR11A1 antibody detected the OR11A1 protein (arrow).

OR11A1 Antibody(C-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR11A1 Antibody(C-term) - References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
Barcellos, L.F., et al. PLoS Genet. 5 (10), E1000696 (2009) :
Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)
Mungall, A.J., et al. Nature 425(6960):805-811(2003)
Volz, A., et al. J. Biol. Chem. 278(22):19691-19701(2003)