10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

VPS25 Antibody(C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19683b

Specification

VPS25 Antibody(C-term) - Product Information

Application	**WB,E**
Primary Accession	Q9BRG1
Other Accession	Q5E9A6, NP_115729.1
Reactivity	**Mouse**
Predicted	**Bovine**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**20748**
Antigen Region	**116-144**

VPS25 Antibody(C-term) - Additional Information

Gene ID 84313

Other Names
Vacuolar protein-sorting-associated protein 25, hVps25, Dermal papilla-derived protein 9, ELL-associated protein of 20 kDa, ESCRT-II complex subunit VPS25, VPS25, DERP9, EAP20

Target/Specificity
This VPS25 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 116-144 amino acids from the C-terminal region of human VPS25.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
VPS25 Antibody(C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

VPS25 Antibody(C-term) - Protein Information

Name VPS25

Synonyms DERP9, EAP20

Function Component of the ESCRT-II complex (endosomal sorting complex required for transport II), which is required for multivesicular body (MVB) formation and sorting of endosomal cargo proteins into MVBs. The MVB pathway mediates delivery of transmembrane proteins into the lumen of the lysosome for degradation. The ESCRT-II complex is probably involved in the recruitment of the ESCRT-III complex. The ESCRT-II complex may also play a role in transcription regulation, possibly via its interaction with ELL. The ESCRT-II complex may be involved in facilitating the budding of certain RNA viruses.

Cellular Location
Cytoplasm. Endosome membrane. Nucleus, nucleoplasm. Note=Distributes diffusely throughout the cytoplasm and nucleoplasm, but exhibits a punctate distribution on coexpression with CHMP6


Tissue Location
Expressed at the mRNA level in kidney, liver, pancreas, and placenta. Lower levels of expression are found in heart, skeletal muscle, brain and lung.

VPS25 Antibody(C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

VPS25 Antibody(C-term) - Images



VPS25 Antibody (C-term) (Cat. #AP19683b) western blot analysis in kidney cell line lysates (35ug/lane).This demonstrates the VPS25 antibody detected the VPS25 protein (arrow).

VPS25 Antibody(C-term) - Background

VPS25, VPS36 (MIM 610903), and SNF8 (MIM 610904) form ESCRT-II (endosomal sorting complex required for transport II), a

complex involved in endocytosis of ubiquitinated membrane proteins. VPS25, VPS36, and SNF8 are also associated in a multiprotein complex with RNA polymerase II elongation factor (ELL; MIM 600284) (Slagsvold et al., 2005 [PubMed 15755741]; Kamura et al., 2001 [PubMed 11278625]).

VPS25 Antibody(C-term) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)
Im, Y.J., et al. Dev. Cell 17(2):234-243(2009)
Lamesch, P., et al. Genomics 89(3):307-315(2007)
Bowers, K., et al. J. Biol. Chem. 281(8):5094-5105(2006)