

OR5AC2 Antibody(C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19866b**Specification**

OR5AC2 Antibody(C-term) - Product Information

Application	WB,E
Primary Accession	O9NZP5
Other Accession	NP_473447.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35304
Antigen Region	246-274

OR5AC2 Antibody(C-term) - Additional Information**Gene ID** 81050**Other Names**

Olfactory receptor 5AC2, HSA1, OR5AC2

Target/Specificity

This OR5AC2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 246-274 amino acids from the C-terminal region of human OR5AC2.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OR5AC2 Antibody(C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OR5AC2 Antibody(C-term) - Protein Information**Name** OR5AC2**Function** Odorant receptor.


Cellular Location

Cell membrane; Multi-pass membrane protein.

OR5AC2 Antibody(C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OR5AC2 Antibody(C-term) - Images

OR5AC2 Antibody (C-term) (Cat. #AP19866b) pre-incubated without(lane 1) and with(lane 2) blocking peptide in A375 cell line lysate. OR5AC2 Antibody (C-term) (arrow) was detected using the purified Pab.

OR5AC2 Antibody(C-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR5AC2 Antibody(C-term) - References

Muzny, D.M., et al. Nature 440(7088):1194-1198(2006)
Rouquier, S., et al. Proc. Natl. Acad. Sci. U.S.A. 97(6):2870-2874(2000)