

SDHB Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19974b**Specification**

SDHB Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P21912
Other Accession	Q007T0 , Q9COA3 , Q3T189 , NP_002991.2
Reactivity	Human
Predicted	Bovine, Mouse, Pig
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	31630
Antigen Region	205-234

SDHB Antibody (C-term) - Additional Information**Gene ID** 6390**Other Names**

Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial, Iron-sulfur subunit of complex II, Ip, SDHB, SDH, SDH1

Target/Specificity

This SDHB antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 205-234 amino acids from the C-terminal region of human SDHB.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SDHB Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SDHB Antibody (C-term) - Protein Information**Name** SDHB

Synonyms SDH, SDH1

Function Iron-sulfur protein (IP) subunit of the succinate dehydrogenase complex (mitochondrial respiratory chain complex II), responsible for transferring electrons from succinate to ubiquinone (coenzyme Q).


Cellular Location

Mitochondrion inner membrane {ECO:0000250|UniProtKB:Q9YHT2}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q9YHT2}; Matrix side {ECO:0000250|UniProtKB:Q9YHT2}

SDHB Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SDHB Antibody (C-term) - Images

SDHB Antibody (C-term) (Cat. #AP19974b) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the SDHB antibody detected the SDHB protein (arrow).

SDHB Antibody (C-term) - Background

Complex II of the respiratory chain, which is specifically involved in the oxidation of succinate, carries electrons from FADH to CoQ. The complex is composed of four nuclear-encoded subunits and is localized in the mitochondrial inner membrane. The iron-sulfur subunit is highly conserved and contains three cysteine-rich clusters which may comprise the iron-sulfur centers of the enzyme. Sporadic and familial mutations in this gene result in paragangliomas and pheochromocytoma, and support a link between mitochondrial dysfunction and tumorigenesis.

SDHB Antibody (C-term) - References

Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)
Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Cerecer-Gil, N.Y., et al. Clin. Cancer Res. 16(16):4148-4154(2010)
Schimke, R.N., et al. Am. J. Med. Genet. A 152A (6), 1531-1535 (2010) :
Hes, F.J., et al. BMC Med. Genet. 11, 92 (2010) :

SDHB Antibody (C-term) - Citations

- [Stable nuclear expression of ATP8 and ATP6 genes rescues a mtDNA Complex V null mutant.](#)
- [The effect of 3-bromopyruvate on human colorectal cancer cells is dependent on glucose concentration but not hexokinase II expression.](#)