10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

RGS2 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20183a

Specification

RGS2 Antibody (N-term) - Product Information

Application	**WB,E**
Primary Accession	P41220
Other Accession	NP_002914.1
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**24382**
Antigen Region	**4-32**

RGS2 Antibody (N-term) - Additional Information

Gene ID 5997

Other Names
Regulator of G-protein signaling 2, RGS2, Cell growth-inhibiting gene 31 protein, G0/G1 switch regulatory protein 8, RGS2, G0S8

Target/Specificity
This RGS2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 4-32 amino acids from the N-terminal region of human RGS2.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
RGS2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

RGS2 Antibody (N-term) - Protein Information

Name RGS2

Synonyms G0S8

Function Regulates G protein-coupled receptor signaling cascades. Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits, thereby driving them into their inactive GDP- bound form (PubMed:11063746, PubMed:19478087). It is involved in the negative regulation of the angiotensin-activated signaling pathway (PubMed:28784619). Plays a role in the regulation of blood pressure in response to signaling via G protein-coupled receptors and GNAQ. Plays a role in regulating the constriction and relaxation of vascular smooth muscle (By similarity). Binds EIF2B5 and blocks its activity, thereby inhibiting the translation of mRNA into protein (PubMed:19736320).

Cellular Location
[Isoform 1]: Cell membrane. Cytoplasm. Nucleus, nucleolus [Isoform 3]: Cell membrane. Cytoplasm Nucleus, nucleolus


Tissue Location
Expressed in acute myelogenous leukemia (AML) and in acute lymphoblastic leukemia (ALL).

RGS2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

RGS2 Antibody (N-term) - Images



RGS2 Antibody (N-term) (Cat. #AP20183a) western blot analysis in HL-60 cell line lysates (35ug/lane).This demonstrates the RGS2 antibody detected the RGS2 protein (arrow).

RGS2 Antibody (N-term) - Background

Regulator of G protein signaling (RGS) family members are regulatory molecules that act as GTPase activating proteins (GAPs) for G alpha subunits of heterotrimeric G proteins. RGS proteins are able to deactivate G protein subunits of the Gi alpha, Go alpha and

Gq alpha subtypes. They drive G proteins into their inactive GDP-bound forms. Regulator of G protein signaling 2 belongs to this family. The protein acts as a mediator of myeloid differentiation and may play a role in leukemogenesis.

RGS2 Antibody (N-term) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Wang, J., et al. Carcinogenesis 31(10):1755-1761(2010)
Miyamoto-Matsubara, M., et al. Ann. N. Y. Acad. Sci. 1200, 112-119 (2010) :
Li, N.F., et al. Clin. Exp. Hypertens. 32(5):256-261(2010)
Johnatty, S.E., et al. PLoS Genet. 6 (7), E1001016 (2010) :