

Neurogenin1 (NeuroG1) Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2022a**Specification**

Neurogenin1 (NeuroG1) Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	O92886
Other Accession	NP_006152
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	25718
Antigen Region	31-60

Neurogenin1 (NeuroG1) Antibody (N-term) - Additional Information**Gene ID** 4762**Other Names**

Neurogenin-1, NGN-1, Class A basic helix-loop-helix protein 6, bHLHa6, Neurogenic basic-helix-loop-helix protein, Neurogenic differentiation factor 3, NeuroD3, NEUROG1, BHLHA6, NEUROD3, NGN, NGN1

Target/Specificity

This Neurogenin1 (NeuroG1) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 31-60 amino acids from the N-terminal region of human Neurogenin1 (NeuroG1).

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Neurogenin1 (NeuroG1) Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Neurogenin1 (NeuroG1) Antibody (N-term) - Protein Information**Name** NEUROG1

Synonyms BHLHA6, NEUROD3, NGN, NGN1

Function Acts as a transcriptional regulator. Involved in the initiation of neuronal differentiation. Activates transcription by binding to the E box (5'-CANNTG-3'). Associates with chromatin to enhancer regulatory elements in genes encoding key transcriptional regulators of neurogenesis (By similarity).

Cellular Location

Nucleus.

Tissue Location


Expression restricted to the embryonic nervous system

Neurogenin1 (NeuroG1) Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Neurogenin1 (NeuroG1) Antibody (N-term) - Images



Anti-NeuroG1 Antibody (A46) at 1:1000 dilution + HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 26 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



The anti-NeuroG1 N-term Pab (Cat. #AP2022b) is used in Western blot to detect NeuroG1 in A375 cell lysate.

Neurogenin1 (NeuroG1) Antibody (N-term) - Background

Basic helix-loop-helix (bHLH) proteins are transcription factors involved in determining cell type during development. NeuroG1 is a bHLH protein with dual cell-fate specification roles. It functions during neurogenesis, and it has also been shown to inhibit the differentiation of neural stem cells into astrocytes. NeuroG1 promotes neurogenesis by functioning as a transcriptional activator, yet it inhibits astrocyte differentiation by compartmentalizing the CREB-binding protein transcription complex away from astrocyte differentiation genes and by inhibiting STAT transcription factors essential for gliogenesis.

Neurogenin1 (NeuroG1) Antibody (N-term) - References

Tamimi, R.M., et al., Genomics 40(2):355-357 (1997).
McCormick, M.B., et al., Mol. Cell. Biol. 16(10):5792-5800 (1996).