

# Mouse Mast1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP20323b

### **Specification**

# Mouse Mast1 Antibody (C-term) - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Isotype	WB,E <u>O9R1L5</u> <u>O810W7</u> Mouse Rat Rabbit Polyclonal Rabbit IgG
Clonality	Polyclonal
Isotype	
Calculated MW	170997
Antigen Region	1438-1464

# Mouse Mast1 Antibody (C-term) - Additional Information

### Gene ID 56527

**Other Names** Microtubule-associated serine/threonine-protein kinase 1, Syntrophin-associated serine/threonine-protein kinase, Mast1 {ECO:0000312|MGI:1861901}

#### Target/Specificity

This Mouse Mast1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1438-1464 amino acids from the C-terminal region of mouse Mast1.

Dilution WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

Mouse Mast1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

# Mouse Mast1 Antibody (C-term) - Protein Information

Name Mast1 {ECO:0000312|MGI:MGI:1861901}



**Function** Microtubule-associated protein essential for correct brain development (PubMed:<u>30449657</u>). Appears to link the dystrophin/utrophin network with microtubule filaments via the syntrophins. Phosphorylation of DMD or UTRN may modulate their affinities for associated proteins.

# **Cellular Location**

Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cytoskeleton. Cell projection, axon. Cell projection, dendrite. Note=Also localized in the soma of neurons (PubMed:30449657). Observed as punctate clusters in the processes of interneurons and along the cell body periphery Colocalizes with syntrophins at the cell membrane

### **Tissue Location**


Expressed in the brain (PubMed:30449657). Expressed in the developing cortical plate, the intermediate zone and corpus callosal fibers that cross the midline (PubMed:30449657). Detected at low levels in the testis, liver and spleen (PubMed:30449657). Expressed in proximity to neuronal nuclei throughout the cortex and cerebellum, and in the vascular endothelium. Also detected in ependymal cells, the choroid plexus, and in developing spermatid acrosomes

# Mouse Mast1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

### Mouse Mast1 Antibody (C-term) - Images



Mouse Mast1 Antibody (C-term) (Cat. #AP20323b) western blot analysis in mouse spleen tissue lysates (35ug/lane). This demonstrates the Mouse Mast1 antibody detected the Mouse Mast1 protein (arrow).

### Mouse Mast1 Antibody (C-term) - Background



Appears to link the dystrophin/utrophin network with microtubule filaments via the syntrophins. Phosphorylation of DMD or UTRN may modulate their affinities for associated proteins.