

ETV7 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20366b

Specification

ETV7 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O9Y603
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	38998
Antigen Region	313-341

ETV7 Antibody (C-term) - Additional Information

Gene ID 51513

Other Names

Transcription factor ETV7, ETS translocation variant 7, ETS-related protein Tel2, Tel-related Ets factor, Transcription factor Tel-2, ETV7, TEL2, TELB, TREF

Target/Specificity

This ETV7 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 313-341 amino acids from the C-terminal region of human ETV7.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ETV7 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ETV7 Antibody (C-term) - Protein Information

Name ETV7

Synonyms TEL2, TELB, TREF

Function Transcriptional repressor; binds to the DNA sequence 5'- CCGGAAGT-3'. Isoform A does not seem to have a repressor activity. Isoform C does not seem to have a repressor activity.

Cellular Location

Nucleus.

Tissue Location


Expressed in hematopoietic tissues.

ETV7 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ETV7 Antibody (C-term) - Images



All lanes : Anti-ETV7 Antibody (C-term) at 1:2000 dilution Lane 1: A431 whole cell lysate Lane 2: Jurkat whole cell lysate Lane 3: K562 whole cell lysate Lane 4: NCI-H292 whole cell lysate Lane 5: U-87 MG whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 39 kDa Blocking/Dilution buffer: 5% NFD/MTBST.

ETV7 Antibody (C-term) - Background

Transcriptional repressor; binds to the DNA sequence 5'-CCGGAAGT-3'. Isoform A does not seem to have a repressor activity. Isoform C does not seem to have a repressor activity.