

### LRRC61 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP20391c

### **Specification**

# LRRC61 Antibody (Center) - Product Information

Application WB.E **Primary Accession 09BV99** Reactivity Human Host Rabbit Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 28014 **Antigen Region** 111-139

### LRRC61 Antibody (Center) - Additional Information

### **Gene ID** 65999

### **Other Names**

Leucine-rich repeat-containing protein 61, LRRC61

# **Target/Specificity**

This LRRC61 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 111-139 amino acids from the Central region of human LRRC61.

#### **Dilution**

WB~~1:1000

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

# Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**


LRRC61 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

### LRRC61 Antibody (Center) - Protein Information

Name LRRC61

# LRRC61 Antibody (Center) - Protocols








Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## LRRC61 Antibody (Center) - Images



Anti-LRRC61 Antibody (Center) at 1:1000 dilution + K562 whole cell lysate Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 28 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

### LRRC61 Antibody (Center) - Background

LRRC61 contains two Leucine-rich repeat; these are short sequence motifs present in a number of proteins with diverse functions and cellular locations. These repeats are usually involved in protein-protein interactions. Each Leucine Rich Repeat is composed of a beta-alpha unit. These units form elongated non-globular structures. Leucine Rich Repeats are often flanked by cysteine rich domains.