10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CCNB2 Antibody (Center S204)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20413c

Specification

CCNB2 Antibody (Center S204) - Product Information

Application	**WB,E**
Primary Accession	O95067
Other Accession	Q4R7A8
Reactivity	**Human**
Predicted	**Monkey**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**45282**
Antigen Region	**182-209**

CCNB2 Antibody (Center S204) - Additional Information

Gene ID 9133

Other Names
G2/mitotic-specific cyclin-B2, CCNB2

Target/Specificity
This CCNB2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 182-209 amino acids from the Central region of human CCNB2.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
CCNB2 Antibody (Center S204) is for research use only and not for use in diagnostic or therapeutic procedures.

CCNB2 Antibody (Center S204) - Protein Information

Name CCNB2

Function Essential for the control of the cell cycle at the G2/M (mitosis) transition.


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CCNB2 Antibody (Center S204) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CCNB2 Antibody (Center S204) - Images



CCNB2 Antibody (Center S204) (Cat. #AP20413c) western blot analysis in U251 cell line lysates (35ug/lane).This demonstrates the CCNB2 antibody detected the CCNB2 protein (arrow).

CCNB2 Antibody (Center S204) - Background

Essential for the control of the cell cycle at the G2/M (mitosis) transition.