10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MORC3 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20433c

Specification

MORC3 Antibody (Center) - Product Information

Application	**WB, IHC-P,E**
Primary Accession	Q14149
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Antigen Region	**634-663**

MORC3 Antibody (Center) - Additional Information

Gene ID 23515

Other Names
MORC family CW-type zinc finger protein 3, Zinc finger CW-type coiled-coil domain protein 3, MORC3, KIAA0136, ZCWCC3

Target/Specificity
This MORC3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 634-663 amino acids from the Central region of human MORC3.

Dilution
WB~~1:1000
IHC-P~~1:100

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
MORC3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

MORC3 Antibody (Center) - Protein Information

Name MORC3 (HGNC:23572)

Function Nuclear matrix protein which forms MORC3-NBs (nuclear bodies) via an ATP-dependent mechanism and plays a role in innate immunity by restricting different viruses through modulation

of the IFN response (PubMed:27440897, PubMed:34759314). Mechanistically, possesses a primary antiviral function through a MORC3-regulated element that activates IFNB1, and this function is guarded by a secondary IFN- repressing function (PubMed:34759314). Sumoylated MORC3-NBs associates with PML-NBs and recruits TP53 and SP100, thus regulating TP53 activity (PubMed:17332504, PubMed:20501696). Binds RNA in vitro (PubMed:11927593). Histone methylation reader which binds to non- methylated (H3K4me0), monomethylated (H3K4me1), dimethylated (H3K4me2) and trimethylated (H3K4me3) 'Lys-4' on histone H3 (PubMed:26933034). The order of binding preference is H3K4me3 > H3K4me2 > H3K4me1 > H3K4me0 (PubMed:26933034).

Cellular Location
Nucleus, nucleoplasm. Nucleus matrix Nucleus, PML body. Chromosome {ECO:0000250|UniProtKB:F7BJB9}. Note=Also found in PML-independent nuclear bodies. Localization to nuclear bodies is ATP-dependent

Tissue Location
Expressed in heart, placenta, skeletal muscle, brain, pancreas, lung, liver, but not kidney

MORC3 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

MORC3 Antibody (Center) - Images



MORC3 Antibody (Center) (Cat. #AP20433c) western blot analysis in HL-60 and U87-MG cell line lysates (35ug/lane).This demonstrates the MORC3 antibody detected the MORC3 protein (arrow).

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Immunohistochemical analysis of paraffin-embedded H. stomach section using MORC3 Antibody (Center)(Cat#AP20433c). AP20433c was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Immunohistochemical analysis of paraffin-embedded H. kideny section using MORC3 Antibody (Center)(Cat#AP20433c). AP20433c was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Formalin-fixed and paraffin-embedded H.placenta tissue reacted with MORC3 Antibody (Center)

(Cat#AP20433c).

MORC3 Antibody (Center) - Background

This gene encodes a protein that localizes to the nuclear matrix. The protein also has RNA binding activity, and has a predicted coiled coil domain.

MORC3 Antibody (Center) - References

Nagase T., et al. DNA Res. 2:167-174(1995).
Ota T., et al. Nat. Genet. 36:40-45(2004).
Hattori M., et al. Nature 405:311-319(2000).
Burkard T.R., et al. BMC Syst. Biol. 5:17-17(2011).