

### **CNO Antibody (Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP20464c

### **Specification**

# **CNO Antibody (Center) - Product Information**

| WB,E          |
|---------------|
| <u>Q9NUP1</u> |
| Human         |
| Rabbit        |
| Polyclonal    |
| Rabbit IgG    |
| 23351         |
| 114-143       |
|               |

## **CNO Antibody (Center) - Additional Information**

#### Gene ID 55330

**Other Names** Biogenesis of lysosome-related organelles complex 1 subunit 4, BLOC-1 subunit 4, Protein cappuccino homolog, BLOC1S4, CNO

### Target/Specificity

This CNO antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 114-143 amino acids from the Central region of human CNO.

Dilution WB~~1:1000

#### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

CNO Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

### **CNO Antibody (Center) - Protein Information**

Name BLOC1S4

Synonyms CNO



**Function** Component of the BLOC-1 complex, a complex that is required for normal biogenesis of lysosome-related organelles (LRO), such as platelet dense granules and melanosomes. In concert with the AP-3 complex, the BLOC-1 complex is required to target membrane protein cargos into vesicles assembled at cell bodies for delivery into neurites and nerve terminals. The BLOC-1 complex, in association with SNARE proteins, is also proposed to be involved in neurite extension. Plays a role in intracellular vesicle trafficking.


Cellular Location Cytoplasm.

## **CNO Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

## CNO Antibody (Center) - Images



CNO Antibody (Center) (Cat. #AP20464c) western blot analysis in Hela cell line lysates (35ug/lane).This demonstrates the CNO antibody detected the CNO protein (arrow).

# CNO Antibody (Center) - Background

The BLOC-1 complex is required for normal biogenesis of lysosome-related organelles, such as platelet dense granules and melanosomes. Plays a role in intracellular vesicle trafficking (By similarity).

# **CNO Antibody (Center) - References**

Ota T., et al. Nat. Genet. 36:40-45(2004).



Hillier L.W., et al. Nature 434:724-731(2005).