

PLN Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20550a

Specification

PLN Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	P26678
Other Accession	P61016 , P61015 , P61013 , P61014 , A4IFH6
Reactivity	Human
Predicted	Bovine, Mouse, Pig, Rabbit, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	6109

PLN Antibody (N-term) - Additional Information

Gene ID 5350

Other Names

Cardiac phospholamban, PLB, PLN, PLB

Target/Specificity

This PLN antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1-22 amino acids from the N-terminal region of human PLN.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PLN Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PLN Antibody (N-term) - Protein Information

Name PLN ([HGNC:9080](#))

Synonyms PLB

Function Reversibly inhibits the activity of ATP2A2 in cardiac sarcoplasmic reticulum by decreasing the apparent affinity of the ATPase for Ca^{2+} (PubMed:[28890335](#)). Modulates the contractility of the heart muscle in response to physiological stimuli via its effects on ATP2A2. Modulates calcium re-uptake during muscle relaxation and plays an important role in calcium homeostasis in the heart muscle. The degree of ATP2A2 inhibition depends on the oligomeric state of PLN. ATP2A2 inhibition is alleviated by PLN phosphorylation. Controls intracellular Ca^{2+} levels in elongated spermatids. May play a role in germ cell differentiation (By similarity).

Cellular Location

Endoplasmic reticulum membrane; Single-pass membrane protein. Sarcoplasmic reticulum membrane; Single-pass membrane protein. Mitochondrion membrane {ECO:0000250|UniProtKB:A4IFH6}; Single-pass membrane protein. Membrane {ECO:0000250|UniProtKB:P61014}; Single-pass membrane protein. Note=Colocalizes with HAX1 at the endoplasmic reticulum (PubMed:17241641). Colocalizes with DMPK at the sarcoplasmic reticulum (PubMed:15598648).

Tissue Location


Heart muscle (at protein level).

PLN Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PLN Antibody (N-term) - Images



Western blot analysis of lysate from U266 cell line, using PLN Antibody (N-term) (Cat. #AP20550a). AP20550a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35 µg per lane.

PLN Antibody (N-term) - Background

Reversibly inhibits the activity of ATP2A2 in cardiac sarcoplasmic reticulum by decreasing the apparent affinity of the ATPase for Ca^{2+} . Modulates the contractility of the heart muscle in response to physiological stimuli via its effects on ATP2A2. Modulates calcium re-uptake during muscle relaxation and plays an important role in calcium homeostasis in the heart muscle. The degree of ATP2A2 inhibition depends on the oligomeric state of PLN. ATP2A2 inhibition is alleviated by PLN phosphorylation.

PLN Antibody (N-term) - References

Fujii J., et al. J. Biol. Chem. 266:11669-11675(1991).
Salvatore C.A., et al. Submitted (MAR-1991) to the EMBL/GenBank/DDBJ databases.
McTiernan C.F., et al. J. Mol. Cell. Cardiol. 31:679-692(1999).
Minamisawa S., et al. Biochem. Biophys. Res. Commun. 304:1-4(2003).
Kaliman P., et al. J. Biol. Chem. 280:8016-8021(2005).