

## **TCL1A Antibody (Center)**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP20567a

## Specification

## **TCL1A Antibody (Center) - Product Information**

Application Primary Accession	<b>WB, FC,E</b> P56279
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	13460

## **TCL1A Antibody (Center) - Additional Information**

#### Gene ID 8115

Other Names T-cell leukemia/lymphoma protein 1A, Oncogene TCL-1, Oncogene TCL1, Protein p14 TCL1, TCL1A, TCL1

# **Target/Specificity** This TCL1A antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 32-66 amino acids from the Central region of human TCL1A.

**Dilution** WB~~1:1000 FC~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

TCL1A Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

### TCL1A Antibody (Center) - Protein Information

Name TCL1A

Synonyms TCL1



**Function** Enhances the phosphorylation and activation of AKT1, AKT2 and AKT3. Promotes nuclear translocation of AKT1. Enhances cell proliferation, stabilizes mitochondrial membrane potential and promotes cell survival.

**Cellular Location** 

Cytoplasm. Nucleus. Microsome. Endoplasmic reticulum. Note=Microsomal fraction


Tissue Location

Restricted in the T-cell lineage to immature thymocytes and activated peripheral lymphocytes. Preferentially expressed early in T- and B-lymphocyte differentiation

# TCL1A Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>
- TCL1A Antibody (Center) Images



Western blot analysis of lysate from Ramos cell line, using TCL1A Antibody (Center)(Cat. #AP20567a). AP20567a was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.





Flow cytometric analysis of Ramos cells using TCL1A Antibody (Center)(green, Cat#AP20567a) compared to an isotype control of rabbit IgG(blue). AP20567a was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody.

## TCL1A Antibody (Center) - Background

Enhances the phosphorylation and activation of AKT1, AKT2 and AKT3. Promotes nuclear translocation of AKT1. Enhances cell proliferation, stabilizes mitochondrial membrane potential and promotes cell survival.

## **TCL1A Antibody (Center) - References**

Virgilio L.,et al.Proc. Natl. Acad. Sci. U.S.A. 91:12530-12534(1994). Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases. Fu T.-B.,et al.Cancer Res. 54:6297-6301(1994). Laine J.,et al.Mol. Cell 6:395-407(2000). Pekarsky Y.,et al.Proc. Natl. Acad. Sci. U.S.A. 97:3028-3033(2000).