10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ARPC2 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20763c

Specification

ARPC2 Antibody (C-term) - Product Information

Application	**WB,E**
Primary Accession	O15144
Other Accession	P85970, Q9CVB6, Q3MHR7, Q0IH88, Q6IRB1
Reactivity	**Human**
Predicted	**Xenopus, Bovine, Mouse, Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**34333**

ARPC2 Antibody (C-term) - Additional Information

Gene ID 10109

Other Names
Actin-related protein 2/3 complex subunit 2, Arp2/3 complex 34 kDa subunit, p34-ARC, ARPC2, ARC34

Target/Specificity
This ARPC2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 278-311 amino acids from the C-terminal region of human ARPC2.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
ARPC2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ARPC2 Antibody (C-term) - Protein Information

Name ARPC2

Synonyms ARC34

Function Actin-binding component of the Arp2/3 complex, a multiprotein complex that mediates actin polymerization upon stimulation by nucleation-promoting factor (NPF) (PubMed:9230079). The Arp2/3 complex mediates the formation of branched actin networks in the cytoplasm, providing the force for cell motility (PubMed:9230079). Seems to contact the mother actin filament (PubMed:9230079). In addition to its role in the cytoplasmic cytoskeleton, the Arp2/3 complex also promotes actin polymerization in the nucleus, thereby regulating gene transcription and repair of damaged DNA (PubMed:29925947). The Arp2/3 complex promotes homologous recombination (HR) repair in response to DNA damage by promoting nuclear actin polymerization, leading to drive motility of double-strand breaks (DSBs) (PubMed:29925947).


Cellular Location
Cytoplasm, cytoskeleton. Cell projection. Synapse, synaptosome {ECO:0000250|UniProtKB:Q9CVB6}. Nucleus

ARPC2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ARPC2 Antibody (C-term) - Images



Western blot analysis of lysate from HT-1080 cell line, using ARPC2 Antibody (C-term)(Cat. #AP20763c). AP20763c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.

ARPC2 Antibody (C-term) - Background

Functions as actin-binding component of the Arp2/3 complex which is involved in regulation of actin polymerization and together with an activating nucleation-promoting factor (NPF) mediates

the formation of branched actin networks. Seems to contact the mother actin filament.

ARPC2 Antibody (C-term) - References

Welch M.D.,et al.J. Cell Biol. 138:375-384(1997).
Couch F.J.,et al.Genomics 36:86-99(1996).
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Gevaert K.,et al.Nat. Biotechnol. 21:566-569(2003).
Zhang C.,et al.Submitted (DEC-1998) to the EMBL/GenBank/DDBJ databases.