

CDKN1A Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20797c**Specification**

CDKN1A Antibody (C-term) - Product Information

Application	WB
Primary Accession	P38936
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	18119

CDKN1A Antibody (C-term) - Additional Information**Gene ID** 1026**Other Names**

Cyclin-dependent kinase inhibitor 1, CDK-interacting protein 1, Melanoma differentiation-associated protein 6, MDA-6, p21, CDKN1A, CAP20, CDKN1, CIP1, MDA6, PIC1, SDI1, WAF1

Target/Specificity

This CDKN1A antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 133-164 amino acids from the C-terminal region of human CDKN1A.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CDKN1A Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CDKN1A Antibody (C-term) - Protein Information**Name** CDKN1A ([HGNC:1784](#))

Function Plays an important role in controlling cell cycle progression and DNA damage-induced G2 arrest (PubMed:[9106657](#)). Involved in p53/TP53 mediated inhibition of cellular proliferation in response to DNA damage. Also involved in p53-independent DNA damage-induced G2 arrest mediated by CREB3L1 in astrocytes and osteoblasts (By similarity). Binds to and inhibits cyclin-dependent kinase activity, preventing phosphorylation of critical cyclin-dependent kinase substrates and blocking cell cycle progression. Functions in the nuclear localization and assembly of cyclin D-CDK4 complex and promotes its kinase activity towards RB1. At higher stoichiometric ratios, inhibits the kinase activity of the cyclin D-CDK4 complex. Inhibits DNA synthesis by DNA polymerase delta by competing with POLD3 for PCNA binding (PubMed:[11595739](#)).

Cellular Location

Cytoplasm. Nucleus

Tissue Location

Expressed in all adult tissues, with 5-fold lower levels observed in the brain

CDKN1A Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CDKN1A Antibody (C-term) - Images**CDKN1A Antibody (C-term) - Background**

May be the important intermediate by which p53/TP53 mediates its role as an inhibitor of cellular proliferation in response to DNA damage. Binds to and inhibits cyclin-dependent kinase activity, preventing phosphorylation of critical cyclin-dependent kinase substrates and blocking cell cycle progression. Functions in the nuclear localization and assembly of cyclin D- CDK4 complex and promotes its kinase activity towards RB1. At higher stoichiometric ratios, inhibits the kinase activity of the cyclin D-CDK4 complex.

CDKN1A Antibody (C-term) - References

Harper J.W., et al. Cell 75:805-816(1993).
El-Deiry W.S., et al. Cell 75:817-825(1993).
Xiong Y., et al. Nature 366:701-704(1993).
Jiang H., et al. Mol. Cell. Differ. 1:285-299(1993).
Jiang H., et al. Oncogene 10:1855-1864(1995).