

DNMT1 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20862a**Specification**

DNMT1 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	P26358
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	183165

DNMT1 Antibody (N-term) - Additional Information**Gene ID** 1786**Other Names**

DNA (cytosine-5)-methyltransferase 1, Dnmt1, CXXC-type zinc finger protein 9, DNA methyltransferase Hsa1, DNA MTase Hsa1, MHsa1, MCMT, DNMT1, AIM, CXXC9, DNMT

Target/Specificity

This DNMT1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 363-396 amino acids from the N-terminal region of human DNMT1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DNMT1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

DNMT1 Antibody (N-term) - Protein Information**Name** DNMT1**Synonyms** AIM, CXXC9, DNMT**Function** Methylates CpG residues. Preferentially methylates hemimethylated DNA. Associates

with DNA replication sites in S phase maintaining the methylation pattern in the newly synthesized strand, that is essential for epigenetic inheritance. Associates with chromatin during G2 and M phases to maintain DNA methylation independently of replication. It is responsible for maintaining methylation patterns established in development. DNA methylation is coordinated with methylation of histones. Mediates transcriptional repression by direct binding to HDAC2. In association with DNMT3B and via the recruitment of CTCFL/BORIS, involved in activation of BAG1 gene expression by modulating dimethylation of promoter histone H3 at H3K4 and H3K9. Probably forms a corepressor complex required for activated KRAS- mediated promoter hypermethylation and transcriptional silencing of tumor suppressor genes (TSGs) or other tumor-related genes in colorectal cancer (CRC) cells (PubMed:[24623306](#)). Also required to maintain a transcriptionally repressive state of genes in undifferentiated embryonic stem cells (ESCs) (PubMed:[24623306](#)). Associates at promoter regions of tumor suppressor genes (TSGs) leading to their gene silencing (PubMed:[24623306](#)). Promotes tumor growth (PubMed:[24623306](#)).

Cellular Location

Nucleus. Note=Localized to the perinucleolar region.

Tissue Location


Ubiquitous; highly expressed in fetal tissues, heart, kidney, placenta, peripheral blood mononuclear cells, and expressed at lower levels in spleen, lung, brain, small intestine, colon, liver, and skeletal muscle. Isoform 2 is less expressed than isoform 1.

DNMT1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DNMT1 Antibody (N-term) - Images



Western blot analysis of lysate from Hela cell line, using DNMT1 Antibody (N-term)(Cat. #AP20862a). AP20862a was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.

DNMT1 Antibody (N-term) - Background

Methylates CpG residues. Preferentially methylates hemimethylated DNA. Associates with DNA replication sites in S phase maintaining the methylation pattern in the newly synthesized strand, that is essential for epigenetic inheritance. Associates with chromatin during G2 and M phases to maintain DNA methylation independently of replication. It is responsible for maintaining methylation patterns established in development. DNA methylation is coordinated with methylation of histones. Mediates transcriptional repression by direct binding to HDAC2. In association with DNMT3B and via the recruitment of CTCFL/BORIS, involved in activation of BAG1 gene expression by modulating dimethylation of promoter histone H3 at H3K4 and H3K9.

DNMT1 Antibody (N-term) - References

Yen R.-W.C.,et al.Nucleic Acids Res. 20:2287-2291(1992).
Yoder J.A.,et al.J. Biol. Chem. 271:31092-31097(1996).
Li L.C.,et al.Submitted (AUG-1999) to the EMBL/GenBank/DDBJ databases.
Grimwood J.,et al.Nature 428:529-535(2004).
Hsu D.-W.,et al.Proc. Natl. Acad. Sci. U.S.A. 96:9751-9756(1999).