10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Mouse Sbk1 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20874b

Specification

Mouse Sbk1 Antibody (N-term) - Product Information

Application	**WB,E**
Primary Accession	Q8QZX0
Other Accession	Q9Z335
Reactivity	**Mouse**
Predicted	**Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Antigen Region	**65-98**

Mouse Sbk1 Antibody (N-term) - Additional Information

Gene ID 104175

Other Names
Serine/threonine-protein kinase SBK1, SH3-binding kinase 1, Sbk1, Sbk

Target/Specificity
This Mouse Sbk1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 65-98 amino acids from the N-terminal region of Mouse Sbk1.

Dilution
WB~~1:500-1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
Mouse Sbk1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Sbk1 Antibody (N-term) - Protein Information

Name Sbk1

Synonyms Sbk

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Function May be involved in signal-transduction pathways related to the control of brain development.


Cellular Location
Cytoplasm.

Mouse Sbk1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Mouse Sbk1 Antibody (N-term) - Images



Western blot analysis of lysates from mouse heart, mouse brain tissue lysate (from left to right), using Sbk1 Antibody (N-term)(Cat. #AP20874b). AP20874b was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

Mouse Sbk1 Antibody (N-term) - Background

May be involved in signal-transduction pathways related to the control of brain development (By similarity).