

## HES5 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP20976a

### Specification

# HES5 Antibody (Center) - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Isotype Calculated MW WB,E <u>O5TA89</u> <u>O03062</u>, <u>P70120</u> Human, Mouse Rat Rabbit Polyclonal Rabbit IgG 18226

## **HES5 Antibody (Center) - Additional Information**

### Gene ID 388585

### **Other Names**

Transcription factor HES-5, Class B basic helix-loop-helix protein 38, bHLHb38, Hairy and enhancer of split 5, HES5, BHLHB38

#### Target/Specificity

This HES5 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 81-115 amino acids from the Central region of human HES5.

Dilution WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

HES5 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

# HES5 Antibody (Center) - Protein Information

### Name HES5

Synonyms BHLHB38



**Function** Transcriptional repressor of genes that require a bHLH protein for their transcription. Plays an important role as neurogenesis negative regulator (By similarity).

Cellular Location Nucleus {ECO:0000255|PROSITE-ProRule:PRU00380, ECO:0000255|PROSITE-ProRule:PRU00981}


**Tissue Location** Expressed in fetal heart and brain tumors.

# HES5 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

## **HES5 Antibody (Center) - Images**



Western blot analysis of lysates from U-87 MG, SK-N-MC cell line, human heart and mouse heart tissue(from left to right), using HES5 Antibody (Center)(Cat. #AP20976a). AP20976a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

# HES5 Antibody (Center) - Background

Transcriptional repressor of genes that require a bHLH protein for their transcription. Plays an important role as neurogenesis negative regulator (By similarity).

# HES5 Antibody (Center) - References

Wistow G., et al. Submitted (NOV-2005) to the EMBL/GenBank/DDBJ databases.



Gregory S.G., et al. Nature 441:315-321(2006). Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases. Katoh M., et al. Int. J. Oncol. 25:529-534(2004).