

WDR82 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20977b**Specification**

WDR82 Antibody (N-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	Q6UXN9
Other Accession	Q8BFQ4 , Q6NV31 , Q5ZMV7 , Q58E77 , Q640J6
Reactivity	Human, Mouse, Zebrafish
Predicted	Xenopus, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	50-85

WDR82 Antibody (N-term) - Additional Information**Gene ID** 80335**Other Names**

WD repeat-containing protein 82, Protein TMEM113, Swd2, WDR82, TMEM113, WDR82A

Target/Specificity

This WDR82 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 50-85 amino acids from the N-terminal region of human WDR82.

DilutionWB~~1:1000
IHC-P~~1:250
FC~~1:25**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

WDR82 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

WDR82 Antibody (N-term) - Protein Information**Name** WDR82 {ECO:0000303|PubMed:17998332, ECO:0000312|HGNC:HGNC:28826}

Function Regulatory component of the SET1/COMPASS complex implicated in the tethering of this complex to transcriptional start sites of active genes (PubMed:[17998332](#), PubMed:[18838538](#), PubMed:[20516061](#)). Facilitates histone H3 'Lys-4' methylation (H3K4me) via recruitment of the SETD1A or SETD1B to the 'Ser-5' phosphorylated C-terminal domain (CTD) of RNA polymerase II large subunit (POLR2A) (PubMed:[17998332](#), PubMed:[18838538](#)). Component of PTW/PP1 phosphatase complex, which plays a role in the control of chromatin structure and cell cycle progression during the transition from mitosis into interphase (PubMed:[20516061](#)). Together with ZC3H4, but independently of the SET1 complex, part of a transcription termination checkpoint that promotes transcription termination of long non-coding RNAs (lncRNAs) (PubMed:[33913806](#), PubMed:[33767452](#)). The transcription termination checkpoint is activated by the inefficiently spliced first exon of lncRNAs and promotes transcription termination of lncRNAs and their subsequent degradation by the exosome (PubMed:[33767452](#)).

Cellular Location


Nucleus. Chromosome {ECO:0000250|UniProtKB:Q8BFQ4}. Note=Associates with chromatin (PubMed:20516061). Recruited at sites of high RNA polymerase II occupancy (By similarity). {ECO:0000250|UniProtKB:Q8BFQ4, ECO:0000269|PubMed:20516061}

WDR82 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


WDR82 Antibody (N-term) - Images



All lanes : Anti-WDR82 Antibody (N-term) at 1:1000 dilution Lane 1: F9 whole cell lysate Lane 2: NIH/3T3 whole cell lysate Lane 3: SH-SY5Y whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 35 kDa Blocking/Dilution buffer: 5% NFD/MTBST.



AP20977b staining WDR82 in human brain tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Samples were incubated with primary antibody (1/250) for 1 hours at room temperature. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Overlay histogram showing HepG2 cells stained with AP20977b (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP20977b, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed (1583138) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.

WDR82 Antibody (N-term) - Background

Regulatory component of the SET1 complex implicated in the tethering of this complex to transcriptional start sites of active genes. Facilitates histone H3 'Lys-4' methylation via recruitment of the SETD1A or SETD1B to the 'Ser-5' phosphorylated C-terminal domain (CTD) of RNA polymerase II large subunit (POLR2A). Component of PTW/PP1 phosphatase complex, which plays a role in the control of chromatin structure and cell cycle progression during the transition from mitosis into interphase.

WDR82 Antibody (N-term) - References

Clark H.F.,et al.Genome Res. 13:2265-2270(2003).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Lee J.-H.,et al.J. Biol. Chem. 280:41725-41731(2005).
Higa L.A.,et al.Nat. Cell Biol. 8:1277-1283(2006).