10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Mouse Lhx1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21022a

Specification

Mouse Lhx1 Antibody (Center) - Product Information

Application	**WB,E**
Primary Accession	P63006
Other Accession	P29674, P63007, P48742, P53411
Reactivity	**Human, Mouse**
Predicted	**Chicken, Rat, Xenopus**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**44780**

Mouse Lhx1 Antibody (Center) - Additional Information

Gene ID 16869

Other Names
LIM/homeobox protein Lhx1, LIM homeobox protein 1, Homeobox protein Lim-1, Lhx1, Lim-1, Lim1

Target/Specificity
This Mouse Lhx1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 170-204 amino acids from the Central region of Mouse Lhx1.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
Mouse Lhx1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Lhx1 Antibody (Center) - Protein Information

Name Lhx1

Synonyms Lim-1, Lim1

Function Potential transcription factor. May play a role in early mesoderm formation and later in lateral mesoderm differentiation and neurogenesis.

Cellular Location
Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108, ECO:0000269|PubMed:8793615}


Tissue Location
In mid to late stage embryos, expressed in a restricted region of mesoderm in the primitive streak. At 7.5 days, expressed in a horseshoe shape at the periphery of the node, as well as along both sides of the adjacent notochord. Also present in presumptive lateral and intermediate mesoderm. Later, expression become progressively restricted to intermediate mesoderm, and the developing excretory system including the pronephric region, mesonephros, nephric duct and metanephros. In the metanephros, strongly expressed in renal vesicles and S-shaped and coma-shaped bodies, as well as in the ureteric bud and its derivatives. Also expressed in the dorsal root ganglia. By stage 10.5, also expressed in regions of the central nervous system in the telencephalon through to the spinal cord. In adults, expressed in the cerebellum/medulla and kidney, and at very low levels in the cerebrum.

Mouse Lhx1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Mouse Lhx1 Antibody (Center) - Images



Western blot analysis of lysates from CEM cell line, mouse testis tissue lysate (from left to right), using Lhx1 Antibody (Center)(Cat. #AP21022a). AP21022a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Mouse Lhx1 Antibody (Center) - Background

Potential transcription factor. May play a role in early mesoderm formation and later in lateral mesoderm differentiation and neurogenesis.

Mouse Lhx1 Antibody (Center) - References

Fujii T.,et al.Dev. Dyn. 199:73-83(1994).
Barnes J.D.,et al.Dev. Biol. 161:168-178(1994).
Li Y.,et al.Mamm. Genome 10:444-446(1999).
Karavanov A.A.,et al.Int. J. Dev. Biol. 40:453-461(1996).
Agulnick A.D.,et al.Nature 384:270-272(1996).