

SOX7 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21030c**Specification**

SOX7 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q9BT81
Other Accession	P40646
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	42197

SOX7 Antibody (C-term) - Additional Information**Gene ID** 83595**Other Names**

Transcription factor SOX-7, SOX7

Target/Specificity

This SOX7 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 319-353 amino acids from the C-terminal region of human SOX7.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SOX7 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SOX7 Antibody (C-term) - Protein Information**Name** SOX7

Function Binds to and activates the CDH5 promoter, hence plays a role in the transcriptional regulation of genes expressed in the hemogenic endothelium and blocks further differentiation

into blood precursors (By similarity). May be required for the survival of both hematopoietic and endothelial precursors during specification (By similarity). Competes with GATA4 for binding and activation of the FGF3 promoter (By similarity). Represses Wnt/beta-catenin-stimulated transcription, probably by targeting CTNNB1 to proteasomal degradation. Binds the DNA sequence 5'-AACAAAT-3'.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00267}. Cytoplasm


Tissue Location

Widely expressed in adult and fetal tissues. Present both in mesenchymal and epithelial cells in some adult tissues, including colon. Tends to be down-regulated in prostate adenocarcinomas and colorectal tumors due to promoter hypermethylation

SOX7 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SOX7 Antibody (C-term) - Images

Western blot analysis of lysates from HCT116 cell line and human heart tissue (from left to right), using SOX7 Antibody (C-term)(Cat. #AP21030c). AP21030c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

SOX7 Antibody (C-term) - Background

Binds to and activates the CDH5 promoter, hence plays a role in the transcriptional regulation of

genes expressed in the hemogenic endothelium and blocks further differentiation into blood precursors (By similarity). May be required for the survival of both hematopoietic and endothelial precursors during specification (By similarity). Competes with GATA4 for binding and activation of the FGF3 promoter (By similarity). Represses Wnt/beta-catenin-stimulated transcription, probably by targeting CTNNB1 to proteasomal degradation. Binds the DNA sequence 5'-AACAAAT-3'.

SOX7 Antibody (C-term) - References

Takash W., et al. Nucleic Acids Res. 29:4274-4283(2001).
Ota T., et al. Nat. Genet. 36:40-45(2004).
Kalnine N., et al. Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Guo L., et al. Mol. Cancer Res. 6:1421-1430(2008).