10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

TMEM160 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21058a

Specification

TMEM160 Antibody (N-term) - Product Information

Application	**WB,E**
Primary Accession	Q9NX00
Other Accession	Q9D938, Q24JY6
Reactivity	**Human, Mouse**
Predicted	**Bovine**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**19658**
Antigen Region	**32-64**

TMEM160 Antibody (N-term) - Additional Information

Gene ID 54958

Other Names
Transmembrane protein 160, TMEM160

Target/Specificity
This TMEM160 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 32-64 amino acids from the N-terminal region of human TMEM160.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
TMEM160 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TMEM160 Antibody (N-term) - Protein Information

Name TMEM160 (HGNC:26042)

Cellular Location

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999


Mitochondrion inner membrane; Multi-pass membrane protein

TMEM160 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

TMEM160 Antibody (N-term) - Images



Western blot analysis of lysates from HT-29 cell line, mouse brain tissue lysate (from left to right), using TMEM160 Antibody (N-term)(Cat. #AP21058a). AP21058a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

TMEM160 Antibody (N-term) - References

Ota T.,et al.Nat. Genet. 36:40-45(2004).
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).